



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C07H 21/02, 19/067, 19/10</b> <b>C07H 19/173, 19/20, A61K 31/70</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/02501</b>  <b>(43) International Publication Date:</b> 3 February 1994 (03.02.94)
<b>(21) International Application Number:</b> PCT/US93/06807  <b>(22) International Filing Date:</b> 20 July 1993 (20.07.93)  <b>(30) Priority data:</b> 07/918,362           23 July 1992 (23.07.92)   US 07/967,267           27 October 1992 (27.10.92)   US 07/968,849           30 October 1992 (30.10.92)   US  <b>(60) Parent Applications or Grants</b> (63) Related by Continuation US                           07/967,267 (CIP) Filed on                   27 October 1992 (27.10.92) US                           07/968,849 (CIP) Filed on                   30 October 1992 (30.10.92) US                           07/918,362 (CIP) Filed on                   23 July 1992 (23.07.92)  <b>(71) Applicant (for all designated States except US):</b> ISIS PHARMACEUTICALS, INC. [US/US]; 2280 Faraday Avenue, Carlsbad, CA 92008 (US).		<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> McGEE, Daniel, Peter, Claude [CA/US]; 7706 Canterbury Ct., Boulder, CO 80301 (US). COOK, Phillip, Dan [US/US]; 7340 Bolero Street, Carlsbad, CA 92009 (US). GUINOSSO, Charles, John [US/US]; 313 Eucino Drive, Vista, CA 92083 (US).  <b>(74) Agents:</b> CALDWELL, John, W. et al.; Woodcock Wash- burn Kurtz Mackiewicz & Norris, One Liberty Place, 46th Floor, Philadelphia, PA 19103-7301 (US).  <b>(81) Designated States:</b> AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> NOVEL 2'-O-ALKYL NUCLEOSIDES AND PHOSPHORAMIDITES PROCESSES FOR THE PREPARATION AND USES THEREOF		
<b>(57) Abstract</b>  <p>Novel 2'-O-alkyl guanosine compounds and analogs thereof are provided. Processes for the preparation of 2'-O-alkyl guanosine and analogs thereof are also provided. Processes for preparing 2'-O-alkylated guanosine, uridine, cytidine, and 2,6-diaminopurine 3'-O-phosphoramidites are also provided. Methods for the use of oligonucleotides comprising 2'-O-alkyl guanosine are also provided.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

- 1 -

**NOVEL 2'-O-ALKYL NUCLEOSIDES AND PHOSPHORAMIDITES  
PROCESSES FOR THE PREPARATION AND USES THEREOF**

**CROSS REFERENCE TO RELATED APPLICATION**

This application is a continuation in part of  
5 application U.S. Serial No. 967,267 filed October 27, 1992  
which is a continuation-in-part of U.S. Application Serial No.  
918,362 filed on July 23, 1992 and Application Serial Number  
US91/00243 filed on January 11, 1991 which is a continuation-  
in-part of Application Serial No. 463,358 filed on January 11,  
10 1990 and Application Serial No. 566,977 filed on August 13,  
1990. These applications are assigned to the assignee of the  
present application and are incorporated by reference herein in  
their entirety.

**FIELD OF INVENTION**

15 This invention is directed to processes for the  
preparation of 2'-O-alkyl uridine and cytidine phosphor-  
amidites. This invention is also directed to processes for the  
preparation of 2'-O-alkyl, 3'-O-alkyl and 2',3'-di-O-alkyl 2,6-  
diaminopurine riboside, 2'-O-alkyl guanosine and 2'-O-alkyl  
20 guanosine analogs, phosphoramidites of these compounds and  
methods of use thereof.

**BACKGROUND OF THE INVENTION**

A number of oligonucleotide analogs have been made.  
One class of oligonucleotides that have been synthesized are  
25 the 2'-O-substituted oligonucleotides. Such oligonucleotides

- 2 -

have certain unique and useful properties. In United States patent application serial no. 814,961, filed December 24, 1991, entitled Gapped 2' Modified Phosphorothioate Oligonucleotides, assigned to the same assignee as this application, the entire contents of which are herein incorporated by reference, 2' substituted nucleotides are introduced within an oligonucleotide to induce increased binding of the oligonucleotide to a complementary target strand while allowing expression of RNase H activity to destroy the targeted strand.

10 In a recent article, Sproat, B.S., Beijer, B. and Iribarren, A., *Nucleic Acids Research*, 1990, 18:41, the authors noted further use of 2'-O-methyl substituted oligonucleotides as "valuable antisense probes for studying pre-mRNA splicing and the structure of spliceosomes".

15 2'-O-methyl and ethyl nucleotides and methods of making the same have been reported by a number of authors. Robins, M.J., Naik, S.R. and Lee, A.S.K., *J. Org. Chem.*, 39:1891 (1974) reported a low yield synthesis of 2'-O- and 3'-O-methyl guanosine via a stannous chloride catalyzed monomethylation by diazomethane. As was later reported by

20 Robins, M.J., Hansske, F. and Bernier, S.E., *Can. J. Chem.*, 59:3360 (1981), "convenient and high yield methods have been devised for synthesis of the 2'-O- and 3'-O-methyl ethers of adenosine, cytidine, and uridine... However, guanosine has presented significant difficulties." In the foregoing paper, the authors reported an improved synthesis of 2'-O and 3'-O-methyl guanosine. The synthesis was improved by effecting the stannous chloride catalyzed diazomethane methylation of 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine (2-aminoadenosine) in place

30 of guanosine. The diamino purine moiety was then reduced to the corresponding guanine moiety with adenosine deaminase. In a further diazotation reaction described by Singer and Kusmierek, *Biochemistry* 15: 5052 (1976), a mixture of 2' and 3'-O-ethyl guanosine was reported to result from the treatment

35 of guanosine with diazoethane. The alkylation also resulted in alkylation of the heterocyclic base. The alkylated product was treated with base to remove the ethyl group from the

- 3 -

heterocyclic base. The resulting product was identified by virtue of having the same UV spectrum as that of guanosine, but a Rf differing from the Rf of guanosine.

A further improvement in the synthesis of 2'-O-methyl nucleosides was reported by Inoue, H., Hayase, Y. Imura, A., Iwai, S., Miura, K. and Ohtsuka, E., *Nucleic Acids Research*, 15:6131 (1987). This method of synthesis was effected utilizing CH<sub>3</sub>I in the presence of Ag<sub>2</sub>O. This method proved useful for all of the common nucleotides with the exception of guanosine. As reported by these authors, guanosine proved refractory to this synthetic method. Thus these authors again had to effect the 2'-O-methylation of guanosine with diazomethane. In order to avoid methylation of the amino functionality of the guanine base moiety, the guanine base moiety was blocked with an isobutyryl group. Additionally, to avoid methyl esterification of the 3'-O functionality of the sugar moiety, a TIPDS (tetraisopropyldisiloxane) blocking group was used to block both the 3' and the 5' hydroxyls of the sugar moiety.

Sproat et al., *supra* and Sproat, B.S., Iribarren, A.M., Garcia, R.G. and Beijer, B., *Nucleic Acids Research*, 19:733 (1991) addressed the synthesis of 2'-O-methyl guanosine (and 2'-O-allyl guanosine). In both of these Sproat et al. publications, the investigators presented a further synthetic pathway to 2'-O-methylguanosine and 2'-O-allylguanosine. They characterized the further pathway with respect to the prior known synthetic methods as "avoids(ing)...the use of the highly toxic and potentially explosive reagent diazomethane" and being "far superior to the use of silver oxide/methyl iodide." This same synthetic method of the Sproat et al. investigators is also published in B.S. Sproat and A.I. Lamond, "2'-O-Methyloligoribonucleotides: synthesis and applications," *Oligonucleotides and Analogues*, ed. F. Eckstein, (IRL Press, 1991) which described syntheses of 2'-O-methylribonucleoside-3'-O-phosphoramidites. The uridine phosphoramidite synthesis described therein requires both base and sugar protection of

- 4 -

the starting nucleoside prior to alkylation. Only after the base and sugar protecting groups are in place on the uridine is it then alkylated. Post alkylation, the base protecting group is removed followed by 5'-O-dimethoxytritylation and phosphitylation. The cytidine phosphoramidite synthesis described by Sproat and Lamond utilizes (and thus requires) the base and sugar blocked 2'-O-methyl uridine analog. This analog is then converted to a blocked cytidine analog, the blocking group is removed from the sugar, the analog is dimethoxytritylated and finally phosphitylated. The guanosine phosphoramidite synthesis taught by Sproat and Lamond starts from a 2-amino-6-chloronucleoside having 3' and 5' sugar hydroxy groups blocked. This nucleoside is converted to a 2,6-dichloro derivative. The dichloro compound is then 2'-O alkylated. Following O-alkylation, the dichloro compound is converted to a diazido intermediate. The diazido intermediate is in turn converted to a diamino intermediate. The diamino intermediate is then deaminated to the guanosine analogue. The 2-amino group of the guanosine analogue is blocked followed by dimethoxytritylation and finally phosphitylation. This guanosine procedure is also published in Sproat, et. al., *Nucleic Acids Research*, 1991 19:733.

The above synthetic procedures involve multiple steps and numerous reagent treatments - 9 different reagent treatments for uridine, 10 for cytidine and 12 for guanosine. For the cytidine and guanosine compounds at least one of the reagents that is required is not readily available and thus is a very expensive reagent.

Other groups have taught the preparation of other 2'-O-alkylated nucleosides. 2'-O-methylthiomethylguanosine, was reported by Hansske, F., Madej, D. and Robins, M.J., *Tetrahedron*, 40:125 (1984). It was produced as a minor by-product of an oxidization step during the conversion of guanosine to 9- $\beta$ -D-arabinofuranosylguanine, i.e. the arabino analogue of guanosine. The addition of the 2'-O-methylthiomethyl moiety is an artifact from the DMSO solvent utilized during the oxidization procedure. The 2'-O-methylthiomethyl

- 5 -

derivative of 2,6-diaminopurine riboside was also reported in the Hansske et al. publication. It was also obtained as an artifact from the DMSO solvent.

In addition, Gladkaya, et al., *Khim. Prir. Soedin.*,  
5 1989, 4, 568 discloses N<sub>1</sub>-methyl-2'-O-(tetrahydropyran-2-yl) and 2'-O-methyl guanosine. Sproat, et al., *Nucleic Acids Research*, 1991, 19, 733 teaches the preparation of 2'-O-allyl-guanosine. Allylation of guanosine required a further synthetic pathway. Iribarren, et al., *Proc. Natl. Acad. Sci.*, 1990, 87,  
10 7747 also studied 2'-O-allyl oligoribonucleotides. Iribarren, et al. incorporated 2'-O-methyl-, 2'-O-allyl-, and 2'-O-dimethylallyl-substituted nucleotides into oligoribonucleotides to study the effect of these RNA analogues on antisense analysis. Iribarren found that 2'-O-allyl containing  
15 oligoribonucleotides are resistant to digestion by either RNA or DNA specific nucleases and slightly more resistant to nucleases with dual RNA/DNA specificity, than 2'-O-methyl oligoribonucleotides. However, Iribarren found that 2'-O-dimethylallyl containing oligoribonucleotides exhibited reduced  
20 hybridization to complementary RNA sequences as compared to 2'-O-methyl oligoribonucleotides. Thus, Iribarren suggested that further attempts to prepare alkylated RNA probes, especially those superior to 2'-allyl cytidine containing oligoribonucleotides should be limited to 2'-O-alkyl groups  
25 containing less than five carbon atoms.

Certain oligonucleotides containing 2'-O-alkyl substituted nucleotides are promising candidates for use as human pharmaceuticals. Those having long chain alkyl groups (i.e. four or more carbon atoms) are particularly useful. For  
30 example, long chain alkyl groups may accomodate functional groups in appropriate orientation with the opposing strand upon strand hybridization. Thus 2'-O- long chain alkyl nucleotides such as 2'-O- long chain alkyl guanosine nucleotides are highly desirable in some cases. For use in large scale therapeutic  
35 testing and eventually for human pharmaceutical use, large amounts of these oligonucleotides must be synthesized. The large amounts of oligonucleotides in turn requires large

- 6 -

amounts of the 2'-O-alkyl nucleoside phosphoramidites used in synthesizing the oligonucleotides. Consideration must therefore be given to both cost and purity of the starting phosphoramidites used in the synthesis of such  
5 oligonucleotides. As a general premise, as the number of synthetic steps increases, the cost of manufacture increases. Further as the number of steps increases, quality control problems escalate. In view of this, it is evident that there is a great need for new and improved procedures for preparing  
10 nucleosides and nucleoside phosphoramidites.

#### OBJECTS OF THE INVENTION

It is an object of this invention to provide methods of synthesis of 2'-O-alkylated nucleosides and nucleoside analogues.

15 It is an object of this invention to provide methods of synthesis of 2'-O- and 3'-O-alkylated 2,6-diaminopurine riboside compounds.

It is an object of this invention to provide new and improved synthetic methods for the preparation of 2'-O-alkyl  
20 nucleoside phosphoramidites.

It is an object of this invention is to provide new and improved syntheses of 2'-O-alkyl guanosine phosphoramidites.

It is an object of this invention is to provide new  
25 and improved syntheses of 2'-O-alkyl cytidine phosphoramidites.

It is an object of this invention is to provide new and improved syntheses of 2'-O-alkyl uridine phosphoramidites.

It is an object of this invention is to provide new and improved syntheses of 2,6-diamino-9-(2'-O-alkyl- $\beta$ -D-ribo-  
30 furanosyl)purine phosphoramidites.

It is an object of this invention is to provide new and improved oligonucleotide syntheses that utilize the improved phosphoramidite syntheses of the invention.


These and other objects will become apparent to  
35 persons of ordinary skill in the art from a review of the present specification and appended claims.



- 7 -

**SUMMARY OF THE INVENTION**


This invention includes compounds having the structure:



I

- 5            wherein X is  $R_1-(R_2)_n$ ;  
               $R_1$  is  $C_3-C_{20}$  alkyl,  $C_4-C_{20}$  alkenyl or  $C_2-C_{20}$  alkynyl;  
               $R_2$  is halogen, hydroxyl, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-  
 10 aralkyl, S-aralkyl, NH-aralkyl, amino, N-phthalimido, imidazole, azido, hydrazino, hydroxylamino, isocyanato, sulf-oxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, poly-amine, polyamide, polyalkylene glycol, polyether, a group that  
 15 enhances the pharmacodynamic properties of oligonucleotides, or a group that enhances the pharmacokinetic properties of oligonucleotides; and n is an integer from 0 to about 6.

In other embodiments of the present invention compounds having the structure:



- 8 -

## II

wherein X is  $R_1-(R_2)_n$ ;

$R_1$  is  $C_3-C_{20}$  alkyl;


$R_2$  is  $NH_2$ , H-imidazole, N-phthalimido;

5 Y is a hydroxyl blocking group;

Z is phosphate or an activated phosphate group;

$Q_1$  and  $Q_2$  independently are H or a guanosine blocking group; and n is an integer from 0 to about 6, are also provided.

10 In still other embodiments of the present invention are provided compounds having the structure:



## III

wherein X is  $R_1-(R_2)_n$ ;

$R_1$  is  $C_3-C_{20}$  alkyl,  $C_4-C_{20}$  alkenyl or  $C_2-C_{20}$  alkynyl;

15  $R_2$  is halogen, hydroxyl, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, imidazole, N-phthalimido, azido, hydrazino, hydroxylamino, isocyanato, sulf-  
 20 oxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, polyamine, polyamide, polyalkylene glycol, polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, and a group that enhances the pharmacokinetic properties of  
 25 oligonucleotides; and n is an integer from 0 to about 6.

- 9 -

This invention also includes processes for the facile preparation of 2'-O and 3'-O-monoalkyl or 2',3'-di-O-alkyl substituted guanosine compounds such as compounds of Formulas I, II and III. Except for preparation with diazomethane, heretofore, direct alkylation of guanosine has proven to be refractory. The present invention provides direct 2' and 3' - O-alkylation of 2,6-diamino-9-(- $\beta$ -D-ribofuranosyl)purine, i.e. 2,6-diaminopurine riboside or 2-aminoadenosine, which can be effected followed by deamination of the 2'-O-alkylated 2,6-diamino purine riboside to the corresponding 2'-O-alkylated guanosine. This alkylation can be practiced, if desired, without the use of blocking groups on either the heterocycle or the sugar moieties of the nucleoside. Further unlike the use of diazomethane, which will only yield the methyl alkylation product, alkylation as practiced in this invention is not limited to just methyl alkylation but is used to yield a plenitude of alkyl substituted guanosine and 2,4-diaminopurine riboside compounds. The necessary compounds used in the invention having the formula R-L wherein R is an alkyl group and L is a leaving group, are either commercially available, are known in the literature or can be prepared by procedures analogous to known literature compounds.

The two step alkylation processes of the invention are further distinguished from the six step procedure of the Sproat et al. investigators. See the above-referenced *Nucleic Acids Research*, 18:41 (1990) and *Nucleic Acids Research*, 19:733 (1991) publications. In those procedures 2-amino-6-chloropurine riboside must first be blocked at both the 3' and 5' positions, converted to the 2,6-dichloro derivative, blocked at the 6 purine position, derivatized to the 2'-O-methyl or 2'-O-allyl derivative, converted to 2,6-diamino derivative, deblocked about the 3' and 5' positions and finally deaminated to the 2'-O-methyl or 2'-O-allyl guanosine product.

In accordance with the processes of this invention, alkylation is effected directly on 2,6-diamino-9-(- $\beta$ -D-ribofuranosyl)purine with an appropriate compound having the formula R-L, wherein R is an alkyl group and L is a leaving

- 10 -

group, in the presence of a base of sufficient strength to effect removal of the proton from the 2' or 3' (or both 2' and 3') hydroxyl of the ribofuranosyl sugar moiety of 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine. Alkylation can be limited to mono  
5 alkylation by limiting the amount of either the R-L group or the base to a stoichiometric (or equivalent) amount. Alternatively dialkylation (on both the 2' and 3' positions) can be practiced by use of an excess R-L group and base to concurrently alkylate both the 2' and the 3' positions.

10 While not wishing to be bound by theory, it has been observed that alkylation predominates at the 2' position compared to the 3' position. Generally a ratio of from about 7:3 to about 8:2 of 2' to 3' alkylation products are obtained (as determined by TLC). For both TLC as well as preparative  
15 scale chromatography, the 2' product generally has a faster Rf than the 3' product. Advantage can be taken of this Rf difference to separate the 2'-O- and 3'-O- products from each other or from 2'-O-,3'-O- dialkylated products. Thus the 2' and 3' alkylation products can be separated by procedures such  
20 as silica gel chromatography if desired.

For alkyl groups that are generally larger than propyl, further advantage can be taken of the rate of deamination of the 2' product verse the 3' product for separation of the 2'-O and 3'-O products. Thus mixtures of 2'-  
25 O and 3'-O alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine are subjected to deamination with adenosine deaminase. The enzymatic deamination of the 2'-O product is more facile than deamination of the 3'-O product. This difference in the rate of deamination allows for separation of the deaminated 2'  
30 product, i.e. the 2'-O-alkylated guanosine, from the slower or non-deaminated 3' product, i.e. the 2,6-diamino-9-(3'-O-alkylated- $\beta$ -D-ribofuranosyl)purine. Additionally procedures such as crystallization has been utilized to further separate a 2' product from the corresponding 3' product by separating  
35 the 2'-O-alkylated diaminopurine riboside product from the corresponding 3'-O-alkylated diaminopurine riboside product.

- 11 -

A preferred base utilized for alkylation is sodium hydride. Other suitable bases may also be utilized, however such bases must have sufficient base strength to remove the proton from the 2' (or 3') hydroxyl moiety of the 2,6-diamino-  
5 purine riboside starting material. While not wishing to be bound by theory, generally any base having a  $pK_a$  about 10  $pK_a$  units greater than the  $pK_a$  of the proton of the 2' hydroxyl moiety of the 2,6-diaminopurine riboside starting material may be used. More specifically, bases having a  $pK_b$  greater than  
10 the  $pK_b$  of sodium hydride may conveniently be selected. Such bases can be selected from compilations of base such as those given in Table 1, page 220 of March, J. *Advanced Organic Chemistry*, Wiley-Interscience, John Wiley & Sons, New York, 1985.

15 The alkylation reactions of the invention typically are conducted in DMF as the solvent. Other suitable solvents include DMSO, N-methyl pyrrolidone and sulfolone.

Preferably, deamination is effected by use of deaminase enzymes. Particularly preferred is adenosine  
20 deaminase. Particularly suitable for use is Adenosine Deaminase Type II available from Sigma Chemical Company, St. Louis, MO. Other deamination reagents may also be employed. The deamination reactions of the invention typically are conducted in a mixture solvent containing an organic solvent  
25 and an aqueous buffer. Suitable for use as the organic solvent are DMSO, N-methyl pyrrolidone and sulfolone. In preferred embodiments of the present invention deamination is achieved using DMSO as the organic solvent. Suitable for use as the aqueous buffer are buffers having a pH compatible to the pH  
30 range of use of the deaminase enzyme. Preferred are phosphate buffers such as sodium phosphate and tris buffers.

In order to enrich the 2' product verse 3' product by elimination of any 3' product, a TIPDS (tetraisopropylsiloxane) protecting group is utilized to protect the 3' and 5' hydroxyl  
35 moieties of the sugar portions of the 2,6-diaminopurine riboside. In the same manner, exclusive 3' product would be obtainable by use of a base stable, non-migratory 2'-O-

- 12 -

protecting group. Such base stable, non-migratory protecting groups include but are not limited to tetrahydropyranyl (THP), 4-methoxytetrahydropyran-4-yl (Mthp), 1-[(2-chloro-4-methyl)phenyl-4-methoxypiperidin-4-yl (Ctmp), triphenylmethyl (trityl), mono-, di- and tri-methoxytrityl and other similar protecting groups.

In accordance with this invention there are also provided improved processes for the preparation of 2'-O-alkylated nucleoside phosphoramidites including 2'-O-alkylated guanosine, cytidine and uridine phosphoramidites.

In accordance with methods of the present invention, preparation of a 2'-O-alkylated guanosine 3'-O-phosphoramidite may comprise the steps of alkylating a 2,6-diamino-9-(ribofuranosyl)purine to form a 2,6-diamino-9-(2'-O-alkylated ribofuranosyl)purine; deaminating said 2,6-diamino-9-(2'-O-alkylated ribofuranosyl)purine to form a 2'-O-alkylated guanosine; blocking the 5'-hydroxyl moiety of said 2'-O-alkylated guanosine; and phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated guanosine.

Further in accordance with the invention there 2'-O-alkylated cytidine 3'-O-phosphoramidite may be prepared by the steps of alkylating an unblocked cytidine to form a 2'-O-alkylated cytidine; blocking the 5'-hydroxyl moiety of said 2'-O-alkylated cytidine; and phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated cytidine.

2'-O-alkylated uridine 3'-O-phosphoramidite may be prepared by processes that include the steps of treating a uridine with a dialkyltin oxide to form a 2',3'-O-dialkylstannylene derivative of uridine; alkylating said stannylene derivative of uridine to form a 2'-O-alkylated uridine; blocking the 5'-hydroxyl moiety of said 2'-O-alkylated uridine; and phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated uridine.

The 3'-O-phosphoramidite of 2'-O-alkyl guanosine and 2,6-diamino-9-(2'-O-alkyl- $\beta$ -D-ribofuranosyl) purine can be

- 13 -

provided in some embodiments of the present invention by reaction of 2-NH<sub>2</sub>, 5'-OH protected 2'-O-alkyl guanosine or 2-NH<sub>2</sub>, 6-NH<sub>2</sub>, and 5'-OH protected 2,6-diamino-9-(2'-O-alkyl-β-D-ribofuranosyl) purine with commercially available reagent known  
5 to those skilled in the art such as 2-cyanoethyl N,N-diisopropylaminochlorophosphine.

2'-O-alkyl guanosine and 2'-O-alkyl-2,6-diaminopurine riboside may be phosphitylated at the 3'-OH to provide phosphoramidites by methods known in the art such as by  
10 protection of the NH<sub>2</sub> moieties (2- or 2- and 6- NH<sub>2</sub>, respectively) and 5'-OH moiety followed by reaction with cyanoethyl N,N-diisopropyl aminochlorophosphine.

Compounds of the present invention such as are provided herein can be incorporated into oligomers by  
15 procedures known to those skilled in the art.

In accordance with methods of the present invention an oligonucleotide that includes at least one 2'-O-alkylated guanosine nucleotide within the oligonucleotide may be prepared by processes comprising the steps of alkylating a 2,6-diamino-  
20 9-(ribofuranosyl)purine to form a 2,6-diamino-9-(2'-O-alkylated ribofuranosyl)purine; deaminating said 2,6-diamino-9-(2'-O-alkylated ribofuranosyl)purine to form a 2'-O-alkylated guanosine; blocking the 5'-hydroxyl moiety of said 2'-O-alkylated guanosine; phosphitylating the 3'-position of said 5'-blocked  
25 2'-O-alkylated guanosine to form a 2'-O-alkylated guanosine 3'-O-phosphoramidite; and coupling, utilizing phosphoramidite coupling conditions, said 2'-O-alkylated guanosine 3'-O-phosphoramidite to a 5'-hydroxyl moiety of an oligonucleotide.

Further in accordance with the invention there are  
30 provided processes for preparing an oligonucleotide that include at least one 2'-O-alkylated cytidine nucleotide within the sequence of the oligonucleotide, the processes comprise the steps of alkylating a cytidine to provide a 2'-O-alkylated cytidine; blocking the 5'-hydroxyl moiety of said 2'-O-  
35 alkylated cytidine; phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated cytidine to form a 2'-O-alkylated

- 14 -

cytidine 3'-O-phosphoramidite; and coupling, utilizing phosphoramidite coupling chemistry, said 2'-O-alkylated cytidine 3'-O-phosphoramidite to a 5'-hydroxyl moiety of an oligonucleotide.


5 Further in accordance with the invention there are provided processes for preparing an oligonucleotide that include at least one 2'-O-alkylated uridine nucleotide within the sequence of the oligonucleotide, the processes comprise the steps of treating uridine with a dialkyltin oxide to form a  
10 2',3'-O-dialkylstannylene derivative of uridine; alkylating said stannylene derivative to provide a 2'-O-alkylated uridine; blocking the 5'-hydroxyl moiety of said 2'-O-alkylated uridine; phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated uridine to form a 2'-O-alkylated uridine 3'-O-  
15 phosphoramidite; and coupling, utilizing phosphoramidite chemistry, said 2'-O-alkylated uridine 3'-O-phosphoramidite to a 5'-hydroxyl moiety of an oligonucleotide.

Further in accordance with the invention there are provided processes for preparing an oligonucleotide that  
20 include at least one 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine nucleotide within the sequence of the oligonucleotide, the processes comprise the steps of alkylating a 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine to provide a 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine; blocking the  
25 5'-hydroxyl moiety of said 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine; phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)-purine to form a 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine 3'-O-phosphoramidite; and coupling, utilizing  
30 phosphoramidite chemistry, said 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine 3'-O-phosphoramidite to a 5'-hydroxyl moiety of an oligonucleotide.

Oligomers of the present invention may contain at least one subunit having the structure:



- 15 -



wherein X is  $R_1-(R_2)_n$ ;


$R_1$  is  $C_3-C_{20}$  alkyl,  $C_4-C_{20}$  alkenyl  
 $C_2-C_{20}$  alkynyl;

- 5  $R_2$  is halogen, hydroxyl, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, imidazole, N-phthalimido, azido, hydrazino, hydroxylamino, isocyanato, sulf-  
 10 oxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, polyamine, polyamide, polyalkylene glycol, polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, or a group that enhances the pharmacokinetic properties of  
 15 oligonucleotides;

$T_3$  and  $T_5$  independently are OH or a further nucleotide or nucleoside of said oligonucleotide or oligonucleoside that is joined to said structure; and

n is an integer from 0 to about 6.

- 20 In still other embodiments of the present invention oligomers of the present invention may contain at least one subunit having the structure:



wherein X is  $R_1-(R_2)_n$ ;

$R_1$  is  $C_1-C_{20}$  alkyl,  $C_2-C_{20}$  alkenyl  $C_2-C_{20}$  alkynyl;

- 25  $R_2$  is halogen, hydroxyl, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl,

- 16 -

S-alkyl, NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, imidazole, N-phthalimido, azido, hydrazino, hydroxylamino, isocyanato, sulf-oxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, poly-amine, polyamide, polyalkylene glycol, polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, or a group that enhances the pharmacokinetic properties of oligonucleotides;

10  $T_3$  and  $T_5$  independently are OH or a further nucleotide or nucleoside of said oligonucleotide or oligonucleoside that is joined to said structure; and

n is an integer from 0 to about 6.

Such oligomers or oligonucleotides may be prepared by  
15 solid state synthesis or by other means known to those skilled in the art.

In the context of this invention, the term "nucleoside" refers to a sugar and a base that are joined together, normally about an "anomeric" carbon on the sugar.  
20 Both  $\alpha$  and  $\beta$  sugars are encompassed by the present invention. In preferred embodiments of the present invention the nucleoside sugar is a pentofuranosyl sugar, however, other sugars might also be utilized such as carbocyclic or 4'-deoxy-4'-thio sugar analogs.

25 In the context of this invention, the term "oligonucleotide" or "oligomer" refers to a polynucleotide formed from naturally occurring bases and furanosyl groups joined by native phosphodiester bonds. Oligonucleotides of the present invention will, of course, comprise at least one 2'-O-alkyl guanosine or derivative thereof. Thus, this term  
30 effectively refers to naturally occurring species or synthetic species formed from naturally occurring subunits or their close homologs. The term "oligonucleotide" or "oligomer" may also refer to moieties which have portions similar to naturally occurring oligonucleotides but which have non-naturally occurring portions. Thus, oligonucleotides may have altered sugars, altered base moieties, or altered inter-sugar linkages.

- 17 -

Exemplary among these are the phosphorothioate and other sulfur-containing species which are known for use in the art. In accordance with some preferred embodiments, at least some of the phosphodiester bonds of the oligonucleotide have been

5 substituted with a structure which functions to enhance the stability of the oligonucleotide or the ability of the oligonucleotide to penetrate into the region of cells where the messenger RNA is located. It is preferred that such

10 substitutions comprise phosphorothioate bonds, phosphotriesters, methyl phosphonate bonds, short chain alkyl or cycloalkyl structures or short chain heteroatomic or heterocyclic structures. Other preferred substitutions are  $\text{CH}_2\text{-NH-O-CH}_2$ ,  $\text{CH}_2\text{-N(CH}_3\text{)-O-CH}_2$ ,  $\text{CH}_2\text{-O-N(CH}_3\text{)-CH}_2$ ,  $\text{CH}_2\text{-N(CH}_3\text{)-N(CH}_3\text{)-CH}_2$  and  $\text{O-N(CH}_3\text{)-CH}_2\text{-CH}_2$  structures where phosphodiester

15 intersugar linkage is replaced by the substitutions. Also preferred are morpholino structures. Summerton, J.E. and Weller, D.D., U.S. 5,034,506 issued July 23, 1991. In other preferred embodiments, such as the protein-nucleic acid (PNA) backbone, the phosphodiester backbone of the oligonucleotide

20 may be replaced with a polyamide backbone, the bases being bound directly or indirectly to the aza nitrogen atoms of the polyamide backbone. P.E. Nielsen, et al., *Science* 1991 254 1497. In accordance with other preferred embodiments, the phosphodiester bonds are substituted with other structures

25 which are, at once, substantially non-ionic and non-chiral, or with structures which are chiral and enantiomerically specific. Persons of ordinary skill in the art will be able to select other linkages for use in practice of the invention.

Oligonucleotides may also include species which

30 include at least some modified base forms. Thus, purines and pyrimidines other than those normally found in nature may be so employed. Suitable bases include, but are not limited to those described in U.S. Patent 3,687,808. Similarly, modifications on the furanosyl portion of the nucleotide subunits, in

35 addition to 2'-O-alkyl modifications of the present invention, may also be effected, as long as the essential tenets of this invention are adhered to. Examples of such modifications are

- 18 -

2'-halogen-substituted nucleotides. Some specific examples of modifications at the 2' position of sugar moieties which are useful in the present invention are OH, SH, SCH<sub>3</sub>, F, OCN, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, S- or N- alkyl; S- or N- alkenyl; SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>; ONO<sub>2</sub>; NO<sub>2</sub>; N<sub>3</sub>; NH<sub>2</sub>; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a conjugate; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide and other substituents having similar properties. Sugar mimetics such as cyclobutyls may also be used in place of the pentofuranosyl group. Oligonucleotides may also comprise other modifications consistent with the spirit of this invention. Such oligonucleotides are best described as being functionally interchangeable with yet structurally distinct from natural oligonucleotides. All such oligonucleotides are comprehended by this invention so long as they effectively function as subunits in the oligonucleotide.

Preferably oligonucleotides of the present invention are from about 6 to about 50 nucleotides in length. In still more preferred embodiments of the present invention oligonucleotides are from about 12 to about 20 nucleotides in length.

Further as used in this invention, the term "alkylating" refers to the addition of an alkyl, alkenyl or alkynyl moiety, preferably an alkyl moiety, to the precursors of the nucleosides phosphoramidites of the invention. Alkylation of the 2' position of the nucleoside sugar links the alkylating moiety to the 2' position of the sugar via an ether linkage.

Preferred alkyl moieties include un-substituted and substituted straight chain C<sub>1</sub>-C<sub>20</sub> alkyl and un-substituted and substituted branch chain C<sub>1</sub>-C<sub>20</sub> alkyl. Preferred alkenyl groups include un-substituted and substituted straight chain C<sub>2</sub>-C<sub>20</sub> alkenyl, and un-substituted and substituted branch chain C<sub>2</sub>-C<sub>20</sub> alkenyl. Preferred alkynyl groups include un-substituted and

- 19 -

substituted straight chain  $C_2$ - $C_{20}$  alkynyl and un-substituted and substituted branch chain  $C_2$ - $C_{20}$  alkynyl. Thus preferred alkylation groups include but are not limited to  $C_1$  to  $C_{20}$  straight or branched chain lower alkyl or substituted lower alkyl,  $C_2$  to  $C_{20}$  straight or branched chain lower alkenyl or substituted lower alkynyl,  $C_2$  to  $C_{20}$  straight or branched chain lower alkynyl or substituted lower alkynyl.

Alkyl groups of the invention include but are not limited to  $C_1$ - $C_{20}$  straight and branched chained alkyls such as methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, octadecyl, nonadecyl and eicosyl, isopropyl, 2-butyl, isobutyl, 2-methylbutyl, isopentyl, 2-methyl-pentyl, 3-methylpentyl, 2-ethylhexyl and 2-propyl-pentyl. Alkenyl groups include, but are not limited to, unsaturated moieties derived from the above alkyl groups including, but not limited to, vinyl, allyl and crotyl. Alkynyl groups include unsaturated moieties having at least one triple bond that are derived from the above alkyl groups including, but not limited to, ethynyl and propargyl.

Substituent groups for the above include but are not necessarily limited to halogen (Cl, Br, F), hydroxyl (OH), thiol (SH), keto (C=O), carboxyl (COOH), nitrate (ONO<sub>2</sub>), nitro (NO<sub>2</sub>), nitroso (NO), nitrile (CN), trifluoromethyl (CF<sub>3</sub>), trifluoromethoxy (OCF<sub>3</sub>), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH<sub>2</sub>), azido (N<sub>3</sub>), hydrazino (NHNH<sub>2</sub>), hydroxylamino (ONH<sub>2</sub>), isocyanato (OCN), sulfide (SO), sulfone (SO<sub>2</sub>), sulfide (S-), disulfide (S-S), silyl, heterocyclic, alicyclic, carbocyclic, intercalators, reporter molecules, conjugates, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligonucleotides, and groups that enhance the pharmacokinetic properties of oligonucleotides. Such compounds include 3-penten-2-one, 3-methyl-2-butanol, 2-cyano-octyl, 3-methoxy-4-heptanal, 3-nitrobutyl, 4-isopropoxydodecyl, 4-azido-2-nitrodecyl, 5-mercaptononyl, 4-amino-1-pentenyl as well as other substituted groups. These

- 20 -

substituted groups can be introduced in a blocked or protected form and later de-blocked to the parent substituted compound. For example, use of the phthalimido group as a blocked form of an amino substitution is illustrated below.

5           Other suitable substituent groups include intercalators, reporter groups, reporter enzymes, and conjugates including cholesterol, phospholipids, biotin, phenanthroline, phenazine, phenanthridine, anthraquinone, acridones, pyrenes, stilbenes, oxazolo-pyridocarbazoles, anthraquinones, phenanthridines, phenazines, azidobenzenes, psoralens, porphyrins, cholic acids, folic acids fluoresceins, rhodamines, coumarins, and dyes; steroids, lipophilic molecules, peptides, protein, vitamins, RNA cleaving complexes, metal chelators, alkylators and cross-linking agents.

15           One particularly preferred substituent group is  $\text{CF}_3$ . Further particularly preferred substituent groups are phthalimido and imidazole. As noted, use of the phthalimido group allows for introduction of a blocked amino functionality on the alkyl group. Utilizing guanosine analogues prepared in accordance with this invention as intermediates in oligonucleotide synthesis, after oligonucleotide synthesis is complete, the phthalimido group is removed yielding an amino functionality tethered to a guanosine nucleotide within the oligonucleotide sequence. Use of an imidazole moiety as a substituent group on the alkyl functionality introduces the suggested nucleic acid cleaving functionality, imidazole, on a

20  
25

30           Aryl groups include but are not limited to phenyl, tolyl, benzyl, naphthyl, anthracyl, phenanthryl, pyrenyl, and xyl. Halogens include fluorine, chlorine and bromine. Suitable heterocyclic groups include but are not limited to imidazole, tetrazole, triazole, pyrrolidine, piperidine, piperazine and morpholine. Amines include amines of all of the above alkyl, alkenyl, alkynyl and aryl groups including primary and secondary amines and "masked amines" such as phthalimide. Amines are also meant to include polyalkylamino compounds and aminoalkylamines such as aminopropylamine and further

- 21 -

heterocyclo-alkylamines such as imidazol-1, 2 or 4-yl-propylamine. RNA cleaving complexes may be, for example, intercalators or groups which bind in the minor groove of RNA. Intercalators are molecules which insert themselves between  
5 neighboring bases of an oligonucleotide. Reporter molecules are molecules which may aid in the identification of a molecule, either visually or otherwise. Cross-linking agents effectively join two groups.

Suitable leaving groups of the present invention  
10 include halides such as chloride, bromide, and iodide, sulfonates such as tosyl, brosyl, nosyl, mesyl and trifyl and oxonium ions. In preferred embodiments of the present invention the leaving group is a halide. Still other suitable leaving groups are well known to those skilled in the art.

15 In accordance with methods of the present invention, the alkylation is preferably conducted in the presence of a base, preferably a metal hydride such as sodium hydride. Alkylation of the 2',3'-O-dialkylstannylene derivative of uridine preferably is performed in the presence of a salt such  
20 as a metal halide. Cesium fluoride and sodium iodide are preferred in some embodiments of the present invention. Additionally, the 5' hydroxyl blocking group is preferably a dimethoxytrityl moiety. The phosphitylating reagent is preferably bis-N,N-diisopropylaminocynoethylphosphite and the  
25 phosphitylating reaction is preferably conducted in the presence of N,N-diisopropylamino-hydrotetrazolide.

In effecting the alkylation of uridine, 2',3'-O-(dibutylstannylene) uridine is alkylated. The dibutylstannylene derivative in turn was prepared in one step from  
30 uridine by reaction with dibutyl tin oxide utilizing the procedure of by Wagner, D., Verheyden, J.P.H. and Moffat, J.G., *J. Org. Chem.* 1974, 39:24. As noted by these authors, 2',3'-di-O-(dibutylstannylene) nucleosides are activated towards alkylation. By using the dibutylstannylene derivative  
35 alkylation of the sugar hydroxyls was effected without concurrent alkylation of the uracil base. The

- 22 -

dibutylstannylene group thus served as a activating group not a blocking group.

For the synthesis of N4-benzoyl-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-methyl cytidine 3'-O- $\beta$ -  
5 cyanoethyl-N,N-diisopropylaminophosphoramidite two methods for the preparation of the intermediate N4-benzoyl-2'-O-methylcytidine are compared. Method A involves blocking of the 3'-5' sites with the TIPS-Cl reagent to allow methylation only on the 2' position. Method B, a preferred method of the  
10 invention, uses a direct methylation of cytidine followed by separation of the resulting mixture of 2' and 3' isomers. The overall yields are comparable. In using Method B, the 2'-O-isomer can be crystallized out from the mixture, filtered and the remaining mother liquors taken through the  
15 dimethoxytritylation step prior to separation of the 2' and 3' isomers or alternately the totality of the alkylated cytidine can be taken through the dimethoxytritylation step with separation of the 2' isomer only effected after this step.

In effecting the alkylation of guanosine, 2',6  
20 diaminopurine is alkylated, for example, by methods described in Attorney Docket No. ISIS-710, filed October 27, 1992.

The amino moiety of the phosphoramidites of the invention can be selected from various amines presently used for such phosphoramidites. Such amines include both aliphatic  
25 and heteroaryl amines as are described in various United States patents, principally those to M. Caruthers and associates. These include United States patents 4,668,777, issued May 26, 1987; 4,458,066, issued Jul. 3, 1984; 4,415,732, issued Nov. 15, 1983; and 4,500,707, issued Feb. 19, 1985, all of which are  
30 herein incorporated by reference. One preferred amino group is diisopropylamino.

In addition to the amino moiety of the phosphoramidite, for phosphodiester and phosphorothioate linkages, an additional phosphorous blocking group is used. One  
35 preferred blocking group is the cyanoethyl group. Other phosphorous blocking groups include methoxy and 2-(methylsulphonyl)ethyl. Additionally an activating agent is



- 23 -

normally used for the phosphoramidite coupling chemistry. One preferred activating agent is N,N-diisopropylaminohydro-tetrazolide. Other suitable moieties for these functions are also disclosed in the above noted patents as well as in United States patent 4,725,677, issued Feb. 16, 1988 and Berner, S., Muhlegger, K., and Seliger, H., *Nucleic Acids Research* 1989, 17:853; Dahl, B.H., Nielsen, J. and Dahl, O., *Nucleic Acids Research* 1987, 15:1729; and Nielson, J. Marugg, J.E., Van Boom, J.H., Honnens, J., Taagaard, M. and Dahl, O., *J. Chem. Research* 1986, 26, all of which are herein incorporated by reference.

For use in phosphorothioate linkage, the Beaucage reagent is described in Beaucage, S.L. and Caruthers, M.H., *Tetrahedron Letters* 1981, 22:1859 as well as in Zon, G. and Stec, J., *Phosphorothioate oligonucleotides: Oligonucleotides and Analogs A Practical Approach*; Eckstein, F. Ed.; IRL Press, Oxford, 1991, which also describes sulfurization by elemental sulfur.

Antisense therapy involves the use of oligonucleotides which are specifically hybridizable to target RNA or DNA. Oligonucleotides of the present invention are preferably specifically hybridizable with a target region. By "specifically hybridizable" herein is meant capable of forming a stable duplex with a target DNA or RNA. Upon binding to, or forming a stable duplex with, the target RNA or DNA, the antisense oligonucleotide can selectively inhibit the genetic expression of these nucleic acids or can induce some other events such as destruction of a targeted RNA or DNA or activation of gene expression. Destruction of targeted RNA can be effected by RNase H activation or by linking strand cleavers to the oligonucleotide. Antisense therapy is known in the art. See for example, PCT/US91/05720 filed December 3, 1991 entitled "Antisense Oligonucleotide Inhibitors of Papillomavirus" and PCT/US91/01327 filed February 25, 1991 entitled "Oligonucleotide Therapies for Modulating the Effects of Herpesvirus".

In some embodiments of the present invention the oligonucleotide portions of compounds of the present invention

- 24 -

are at least 60% complementary to a target sequence. In preferred embodiments of the present invention the oligonucleotide portions of compounds of the present invention are at least 80% complementary to a target sequence. 100%  
5 complementarity of the oligonucleotide portions of compounds of the present invention to a target sequence is most preferred. In preferred embodiments of the present invention, the oligonucleotide portions may be specifically hybridizable with DNA or RNA from *Candida*, papilloma virus, Epstein Barr virus,  
10 rhinovirus, hepatitis, human immunodeficiency virus, herpes simplex virus, influenza virus and cytomegalovirus.

2'-O-alkyl guanosine containing oligonucleotides and 2,6-diamino purine containing oligonucleotides of the present invention may be used to modulate the production of protein by  
15 contacting a selected sequence of RNA or DNA coding for a selected protein with an 2'-O-alkyl guanosine or 2,6-diamino purine containing oligonucleotide of the present invention having a sequence of nucleotide bases specifically hybridizable with said selected sequence of RNA or DNA coding for said  
20 protein.

The oligonucleotides of the present invention can be used in diagnostics, therapeutics and as research reagents. For therapeutic use, an animal having a disease characterized by the undesired production of a protein is contacted with an  
25 oligonucleotide of the present invention having a sequence of nucleotide bases specifically hybridizable with a selected sequence of RNA or DNA coding for said protein.

#### EXAMPLES

The following examples illustrate the invention,  
30 however, they are not intended as being limiting. In various examples the nomenclature 4,4'-dimethoxytriphenylmethyl and dimethoxytrityl are used interchangeably to reference the DMT blocking group positioned on the 5'-hydroxyl moiety of the various nucleoside and nucleotides of the invention.

35 NMR spectra were obtained with the following instruments: <sup>1</sup>H-NMR: Varian Gemini-200 (199.975 MHz), <sup>13</sup>C-NMR:

- 25 -

Varian Gemini-200 (50.289 MHz). NMR spectra were recorded using either deuteriochloroform (tetramethylsilane as internal standard) or dimethylsulfoxide- $d_6$  as solvent. The following abbreviations were used to designate the multiplicity of individual signals: s = singlet, d = doublet, t = triplet, q = quartet, ABq = ab quartet, m = multiplet, dd = doublet of doublets, br s = broad singlet. Mass spectra were acquired on a VG 70-SEQ instrument (VG Analytical (Fisons)), using fast atom bombardment ionization (7 kV Xe atoms). Solvent ratios for column chromatography are given as volume/volume. Evaporations of solvents were performed *in vacuo* (60 torr) at 30°C unless otherwise specified. Melting points are reported uncorrected.

**EXAMPLE 1****2,6-Diamino-9-( $\beta$ -D-ribofuranosyl)purine**

In accordance with modifications of the procedures described in Robins, M.J., Hanske, F. and Beriner, S.E., *Can. J. Chem.*, 59:3360 (1981), guanosine hydrate (49 g, Aldrich Chemical Co.), toluene (200 ml), hexamethyldisilazane (160 ml, 4.3 eq) and trifluoromethanesulfonic acid (3.7 ml) were loaded in a stainless steel Parr bomb. The bomb was sealed and heated approximately 1/3 submerged in an oil bath at 170° C for 5 days. The bomb was cooled in a dry ice acetone bath and opened. The contents were transferred to a 2 liter round bottom flask using methanol (MeOH) and the solvent evaporated on a Buchii evaporator. 1:1 H<sub>2</sub>O/MeOH (600 ml) was added to the residue and the resulting brown suspension was refluxed 4-5 hr. The resulting suspension was evaporated on the Buchii evaporator to remove the methanol ( $\approx$  1/2 volume). Additional H<sub>2</sub>O ( $\approx$ 300 ml) was added and the mixture was heated, treated with charcoal and filtered through a Celite filter pad. Upon cooling, a crystalline solid formed. The solid was isolated by filtration, washed with H<sub>2</sub>O and dried under high vacuum at 90° C to yield the product (43.7 g, 89% yield) as a tan solid.

- 26 -

UV and NMR spectra of this compound compared to literature values.

This variation of the procedures of Robins, et al. *supra*, eliminated the need to utilize liquid ammonia in the reaction mixture since the ammonia molecule is generated *in situ* from the silazane reagent and the water of hydration of the guanosine hydrate starting material. Further, the use of chlorotrimethylsilane was not necessary nor was it necessary to conduct the reaction under anhydrous conditions, do a preliminary evaporation, or open and re-seal the Parr bomb under a dry nitrogen atmosphere.

#### EXAMPLE 2

2,6-Diamino-9-(2'-O-propyl- $\beta$ -D-ribofuranosyl)purine & 2,6-Diamino-9-(3-O-propyl- $\beta$ -D-ribofuranosyl)purine

Sodium hydride (NaH) (2.1 g) was added to 2,6-diamino-9-( $\beta$ -D-ribofuranosyl) purine (10.5 g) in dry dimethylformamide (DMF) (150 ml). After stirring for 10 min, iodopropane (6 ml) was added. The solution was stirred for 45 min at room temperature followed by the addition of a further aliquot of NaH (600 mg). The reaction mixture was stirred overnight and then quenched by the addition of ethanol (EtOH) (5 ml). The reaction mixture was evaporated *in vacuo*, the residue suspended in 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub> and purified by silica gel chromatography (300 g) using 5  $\rightarrow$  10% MeOH/CH<sub>2</sub>Cl<sub>2</sub> as the eluent. The 2',3'-di-O-propyl product eluted first followed by the 2'-O-propyl product and then the 3'-O-propyl product. The 2'-O-propyl product containing fractions were pooled and the solvent stripped to yield a crude foam. The foam was crystallized from H<sub>2</sub>O (40 ml), washed with cold H<sub>2</sub>O and dried to yield 2.9 g of the 2'-O-propyl compound. The mother liquor was evaporated, re-chromatographed and crystallized to yield an additional 2.4 g of the 2'-O-propyl compound. The second mother liquor was evaporated to yield 4 g of a mixture of 2' and 3'-O-propyl compounds as an oil. Fractions containing the 3'-O-propyl product as the major product were evaporated and residue

- 27 -

crystallized from water. (See Example 17 below for isolation and characterization of the 2',3'-di-O-propyl compound).

**2,6-Diamino-9-(2'-O-propyl- $\beta$ -D-ribofuranosyl)purine**

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  0.76 (t, 3, CH<sub>3</sub>), 1.4 (tq, 2, CH<sub>2</sub>), 3.3 (m, 1, H-5'' + HDO), 3.65-3.45 (m, 3, H-5', O-CH<sub>2</sub>), 3.9 (m, 1), 4.25 (br m, 1), 4.38 (dd, 1), 5.1 (br d, 1 3'-OH), 5.45 (br t, 1, 5'-OH), 5.75 (br s, 2, 6-NH<sub>2</sub>), 5.83 (d, 1, H-1'), 6.77 (br s, 2, 2-NH<sub>2</sub>) and 7.95 (s, 1, H-8). Anal. Calcd. for C<sub>13</sub>H<sub>20</sub>N<sub>6</sub>O<sub>4</sub>·½H<sub>2</sub>O: C, 46.91; H, 6.2; N, 25.25. Found: C, 47.09; H, 6.37; N, 25.33.

**2,6-Diamino-9-(3'-O-propyl- $\beta$ -D-ribofuranosyl)purine**

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  0.75 (t, 3, CH<sub>3</sub>), 1.4 (tq, 2, CH<sub>2</sub>), 3.27-3.5 (ABX 2, O-CH<sub>2</sub>-), 3.5 and 3.6 (ABX, 2, H-5'), 3.9 (m, 1), 4.22 (m, 1), 4.35 (m, 1), 5.1 (br d, 1, 2'-OH), 5.45 (br t, 1, 5'-OH), 5.75 (br s, 2, 6-NH<sub>2</sub>), 5.8 (d, 1, H-1'), 6.8 (br s, 2CH<sub>2</sub>, 2-H<sub>2</sub>) and 7.95 (s, 1, H-8).

**EXAMPLE 3**

**2'-O-Propylguanosine**

A mixture of 2,6-Diamino-9-(2'-O-propyl- $\beta$ -D-ribofuranosyl) purine and 2,6-Diamino-9-(3'-O-propyl- $\beta$ -D-ribofuranosyl) purine (4.6 gm) and adenosine deaminase (200 mg, Sigma Chemicals Type II) were stirred at room temperature overnight in 0.1 M tris buffer (150 ml, pH 7.4), DMSO (100 ml) and 0.1 M sodium phosphate buffer (10 ml). A further aliquot of adenosine deaminase (140 mg) in 0.1 M phosphate buffer (30 ml) and DMSO (20 ml) was added and the reaction stirred an addition 24 hrs. The solvent was evaporated in vacuo and the residue flash chromatographed on silica gel utilizing 5 → 20% MeOH/CH<sub>2</sub>Cl<sub>2</sub>. Product-containing fractions were evaporated in vacuo and the residue crystallized from H<sub>2</sub>O to yield 2.6 gm of product. m.p. dec > 270° C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  0.75 (t, 3, CH<sub>3</sub>), 1.42 (tq, 2, CH<sub>2</sub>), 3.3-3.6 (m, 4, H-5', O-CH<sub>2</sub>), 3.85 (m, 1), 4.2 (m, 1), 4.23 (m, 1), 5.10 (t, 1, 5'-OH), 5.13 (d, 1, 3'-OH), 5.75 (d, 1, H-1'), 6.45 (br s, 2, NH<sub>2</sub>), 7.95 (s, 1, H-8) and 10.67 (br s, 1, NH). Anal. Calcd. for C<sub>13</sub>H<sub>19</sub>N<sub>5</sub>O<sub>5</sub>: C, 47.99; H, 5.89; N, 21.53. Found: C, 47.90, H, 5.85; N, 21.44.

- 28 -

**EXAMPLE 4****N2-Isobutyryl-2'-O-propylguanosine**

2'-O-Propylguanosine (3.6 gm) in pyridine (50 ml) was cooled in an ice bath and trimethylsilyl chloride (8.4 ml, 6 eq.) was added. The reaction mixture was stirred for 30 min and isobutyryl chloride (5.8 ml, 5 eq.) was added. The solution was stirred for 4 hours during which it was allowed to warm to room temperature. The solution was cooled, H<sub>2</sub>O added (10 ml) and the solution was stirred for an additional 30 mins. Concentrated NH<sub>4</sub>OH (10 ml) was added and the solution evaporated in vacuo. The residue was purified by silica gel chromatography using 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to elute the product. Product-containing fractions were evaporated to yield 2.5 g of product as a foam. An analytical sample was re-chromatographed on silica and eluted with CH<sub>2</sub>Cl<sub>2</sub> → 6% MeOH/CH<sub>2</sub>Cl<sub>2</sub>. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.75 (t, 3, CH<sub>3</sub>), 1.13 [d, 6, CH(CH<sub>3</sub>)<sub>2</sub>], 1.4 (m, 2, CH<sub>2</sub>), 2.75 [m, 1, CH(CH<sub>3</sub>)<sub>2</sub>], 3.52 (m, 6, OCH<sub>2</sub>), 3.36 and 3.6 (ABX, 2, H-5'), 3.95 (m, 1), 4.26 (m, 1), 4.33 (m, 1), 5.07 (t, 1, 5'-OH), 5.18 (d, 1, 3'-OH), 5.9 (d, 1, H-1'), 8.25 (s, 1, H-8), 11.65 (br s, 1, NH) and 12.1 (br s, 1, NH). Anal. Calcd. for C<sub>17</sub>H<sub>25</sub>N<sub>5</sub>O<sub>6</sub>·½H<sub>2</sub>O: C, 50.49; H, 6.48; N, 17.32. Found: C, 50.81; H, 6.62; N, 17.04.

**EXAMPLE 5****N2-Isobutyryl-5'-dimethoxytrityl-2'-O-propylguanosine**

N2-Isobutyryl-2'-O-propylguanosine (2.64 g) was co-evaporated with pyridine and then solubilized in pyridine (180 ml). Dimethoxytrityl chloride (2.4 g, 1.1 eq) and dimethylaminopyridine (50mg) were added with stirring at room temperature. The reaction mixture was stirred overnight and evaporated in vacuo. The residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub>/ 2x dil Na<sub>2</sub>CO<sub>3</sub>. The organic phase was dried (MgSO<sub>4</sub>) and evaporated. The residue was purified by silica gel chromatography (1:1 EtOAc/Hex → 5% MeOH/EtOAc, 1% TEA) to yield 4.1 g of product. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.78 (t, 3, CH<sub>3</sub>), 1.12 [d, 6, CH(CH<sub>3</sub>)<sub>2</sub>], 1.46 (m, 2, CH<sub>2</sub>), 2.75 [m, 1, CH(CH<sub>3</sub>)<sub>2</sub>], 3.35 and 3.55 (ABX, 2, H-5'), 3.73 (s, 6, OCH<sub>2</sub>), 4.0 (m, 1), 4.3 (m, 1),

- 29 -

4.4 (m, 1), 5.18 (d, 1, 3'-OH), 5.93 (d, 1, H-1'), 6.8, 7.2, 7.36 (m, 13, DMTr), 8.13 (s, 1, H-8), 11.63 (br s, 1, NH) and 12.1 (br s, 1, NH). Anal. Calcd. for  $C_{38}H_{42}N_5O_8 \cdot H_2O$ : C, 63.83; H, 6.20; N, 9.80. Found: C, 64.22; H, 6.35; N, 9.55.

## 5 EXAMPLE 6

### N2-Isobutyryl-5'-dimethoxytrityl-2'-O-propylguanosine 3'-β-cyanoethyl-N,N-diisopropylphosphoramidite

A  $CH_2Cl_2$  solution of N2-Isobutyryl-5'-dimethoxytrityl-2'-O-propylguanosine (4.1 g), bis-(N,N-diisopropylamino)-2-cyanoethylphosphite (3.7 ml, 2 eq) and N,N-diisopropylammonium tetrazolide (0.5 g, 0.5 eq) was stirred at room temperature overnight. The solution was partitioned against dil.  $Na_2CO_3$  and then dil.  $Na_2CO_3/NaCl$  and dried over  $MgSO_4$ . The solvent was evaporated in vacuo and the residue was purified by silica gel chromatography (120 g, 1%TEA in EtOAc) to yield 5.2 g of product as a foam.  $^{31}P$  NMR ( $CDCl_3$ )  $\delta$  150.5, 150.8.

## EXAMPLE 7

### 2,6-Diamino-9-(2'-O-pentyl-β-D-ribofuranosyl)purine & 2,6-Diamino-9-(3'-O-pentyl-β-D-ribofuranosyl)purine

2,6-Diamino-9-(β-D-ribofuranosyl)purine (10 g) was treated with sodium hydride (1.7 g, 1.2 eq) and bromopentane (5.3 ml, 1.2 eq) in DMF (90 ml) as per the procedure of Example 2. Silica gel chromatography yielded three components. The first eluted component (not characterized but believed to be the 2,3-di-(O-pentyl) compound was isolated as an oil (700 mg). The next component isolated as a foam (3.3 g) was crystallized from MeOH to yield 2.8 g of 2,6-diamino-9-(2'-O-pentyl-β-D-ribofuranosyl)purine. The third component isolated as a solid (200 mg) was crystallized from MeOH to yield 80 mg of 2,6-diamino-9-(3'-O-pentyl-β-D-ribofuranosyl)purine. Fractions containing mixtures of the first and second components were evaporated and the residue crystallized from MeOH to yield a further 900 mg of the 2-O-pentyl compound. Further fraction yielded 1.2 g of a mixture of the 2'-O-pentyl and 3'-O-pentyl compounds.

- 30 -

**2,6-Diamino-9-(2'-O-pentyl-β-D-ribofuranosyl)purine**

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.75 (t, 3, CH<sub>3</sub>), 1.16 (m, 4, CH<sub>2</sub>), 1.39 (m, 2, CH<sub>2</sub>), 3.53 (m, 2, CH<sub>2</sub>), 3.3 and 3.6 (ABX, 2, H-5'), 3.93 (br s, 1), 4.23 (m, 1), 4.38 (m, 1), 5.1 (d, 1 3'-OH), 5.5 (t, 1, 5'-OH), 5.75 (br s, 2, 6-NH<sub>2</sub>), 5.82 (d, 1, H-1'), 6.8 (br s, 2, 2-NH<sub>2</sub>) and 7.93 (s, 1, H-8).

**2,6-Diamino-9-(3'-O-pentyl-β-D-ribofuranosyl)purine**

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.87 (t, 3, CH<sub>3</sub>), 1.3 (m, 4, CH<sub>2</sub>), 1.55 (m, 2, CH<sub>2</sub>), 3.5 (m, 2, O-CH<sub>2</sub>-), 3.6 (m, 2, H-5'), 3.86 (m, 1), 3.95 (m, 1), 4.6 (m, 1), 5.32 (br d, 1 2'-OH), 5.46 (br t, 1, 5'-OH), 5.70 (d, 1, H-1'), 5.75 (br s, 2, 6-NH<sub>2</sub>), 6.76 (br s, 2, 2-NH<sub>2</sub>) and 7.93 (s, 1, H-8).

**EXAMPLE 8****2'-O-Pentylguanosine**

2,6-diamino-9-(2'-O-pentyl-β-D-ribofuranosyl)purine (1.9 g) in 0.1 M sodium phosphate buffer (50 ml, pH 6.0) and DMSO (25 ml) was treated with adenosine deaminase (added in two aliquots - first aliquot 50 mg, second aliquot 80 mg) at 35° C as per the procedure of Example 3 to yield 1.4 g of product.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.8 (t, 3, CH<sub>3</sub>), 1.16 (m, 4, 2x CH<sub>2</sub>), 1.4 (m, 2, CH<sub>2</sub>), 3.38, 3.6 (m, 4, OCH<sub>2</sub>, H-5'), 3.93 (s, 1, H-4'), 4.28 (m, 2, H-2', H-3'), 5.17 (br, 2, 5', 3'-OH), 5.8 (d, 1, H-1'), 6.53 (br s, 2, NH<sub>2</sub>), 8.0 (s, 1, H-8) and 10.68 (br, 1, NH).

**EXAMPLE 9****25 N2-Isobutyryl-2'-O-pentylguanosine**

2'-O-pentylguanosine (2.3 g) in pyridine (35 ml) was treated with trimethylsilyl chloride (4.15 ml, 5 eq) and isobutyryl chloride (3.4 ml, 5 eq) as per the procedure of Example 4 to yield the product as a foam (2.3 g). An analytical sample was crystallized from EtOAc/Hex. m.p. 178-180° C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.75 (t, 3, CH<sub>3</sub>), 1.1 [m, 10, 2x CH<sub>2</sub>, CH(CH<sub>3</sub>)<sub>2</sub>], 1.4 (m, 2, CH<sub>2</sub>), 2.74 [m, 1, CH(CH<sub>3</sub>)<sub>2</sub>], 3.56 (m, 4, OCH<sub>2</sub>, H-5'), 3.93 (m, 1, H-4'), 4.25 (m, 1), 4.34 (m, 1), 5.05 (t, 1, 5'-OH), 5.17 (d, 1, 3'-OH), 5.88 (d, 1, H-1'), 8.27 (s, 1, H-8), 11.65 (br s, 1, NH) and 12.05 (br s, 1, NH). Anal.



- 31 -

Calcd. for  $C_{19}H_{29}N_5O_6$ : C, 53.89; H, 6.90; N, 16.54. Found: 53.75; H, 6.92; N, 16.40

**EXAMPLE 10****N2-Isobutyryl-5'-dimethoxytrityl-2'-O-pentylguanosine**

- 5 N2-Isobutyryl-2'-O-pentylguanosine (2.3 g) was treated with dimethoxytrityl chloride (1.7 g, 1.1 eq), and dimethylaminopyridine (100 mg as a catalyst) in pyridine (50 ml) as per the procedure of Example 5 to yield the product as a foam (2.9 g).  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  0.83 (t, 3,  $CH_3$ ), 1.2 [m, 10, 2x  $CH_2$ ,  
10  $CH(CH_3)_2$ ], 1.48 (m, 2,  $CH_2$ ), 2.78 [m, 1,  $CH(CH_3)_2$ ], 3.4, 3.6 (m, 4,  $OCH_2$ , H-5'), 3.75 (s, 6,  $OCH_3$ ), 4.07 (m, 1), 4.27 (m, 1), 4.42 (m, 1), 5.2 (br d, 1, 3'-OH), 5.95 (d, 1, H-1'), 6.85, 7.25, 7.38 (m, 13, DMTr), 8.15 (s, 1, H-8), 11.67 (br s, 1, NH) and 12.1 (br s, 1, NH). Anal. Calcd. for  $C_{40}H_{47}N_5O_8 \cdot \frac{1}{2}H_2O$ : C, 65.38; H, 6.58; N, 9.53. Found: C, 65.37; H, 6.59; N, 9.39.

**EXAMPLE 11****N2-Isobutyryl-5'-dimethoxytrityl-2'-O-pentylguanosine 3'- $\beta$ -cyanoethyl-N,N-diisopropylphosphoramidite**

- 20 N2-Isobutyryl-5'-dimethoxytrityl-2'-O-pentyl-guanosine (1.7 g) was treated with bis-(N,N-diisopropylamino)-2-cyanoethyl-phosphite (1.48 g) and N,N-diisopropylammonium tetrazolide (200 mg) as per the procedure of Example 6 to yield the product (1.4 g).  $^{31}P$  NMR ( $CDCl_3$ )  $\delta$  150.5, 150.85.

**25 EXAMPLE 12****2,6-Diamino-9-(2'-O-nonyl- $\beta$ -D-ribofuranosyl)purine**

- 2,6-Diamino-9-( $\beta$ -D-ribofuranosyl)purine (50 g, 180 mmol) was treated with sodium hydride (8.8 g, 220 mmol) and bromononane (59 g, 54.4 ml, 285 mmol) in DMF (700 ml) as per  
30 the procedure of Example 2 (the diamino compound in DMF was cooled in an ice bath during the addition of NaH) to yield 83 g of crude product. 50 g of crude product was purified by silica gel chromatography. Fraction containing 2'-O-nonyl and 3'-O-nonyl product were combined to give a 77:23 mixture (29 g)

- 32 -

of the 2' and 3' product. Pure 2'-O-nonyl product is obtained by chromatography.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  0.95 (t, 3,  $\text{CH}_3$ ); 1.17 [m, 12,  $\text{O}-\text{CH}_2-\text{CH}_2-(\text{CH}_2)_6$ ]; 1.42 [m, 2,  $\text{O}-\text{CH}_2\text{CH}_2(\text{CH}_2)_6$ ]; 3.27-3.70 (m, 2,  $\text{H}-5'$ ); 3.50-3.70 [m, 2,  $\text{O}-\text{CH}_2(\text{CH}_2)_7$ ]; 3.95 (m, 1,  $\text{H}-4'$ ), 4.24 (m, 1,  $\text{H}-3'$ ); 4.40 (m, 1,  $\text{H}-2'$ ); 5.10 (d, 1,  $3'-\text{OH}$ ,  $J=5$  Hz); 5.50 (t, 1,  $5'-\text{OH}$ ,  $J=6$  Hz); 5.76 (s, 2,  $2-\text{NH}_2$ ); 5.83 (d, 1,  $\text{H}-1'$ ,  $J=6.0$  Hz); 6.81 (s, 2,  $6-\text{NH}_2$ ); and 7.96 (s, 1,  $8-\text{H}$ ).

**EXAMPLE 13****2'-O-Nonylguanosine**

10 A mixture of 2,6-diamino-9-(2'-O-nonyl- $\beta$ -D-ribofuranosyl)purine and 2,6-diamino-9-(3'-O-nonyl- $\beta$ -D-ribofuranosyl)purine ( $\approx$  80:20 mixture, 29 g) in 0.1 M sodium phosphate buffer (50 ml, pH 7.4), 0.1 M tris buffer (1800 ml, pH 7.4) and DMSO (1080 ml) was treated with adenosine deaminase  
15 (1.6 g) as per the procedure of Example 3 to yield 60 g of product as an oil. An analytical product was purified by silica gel chromatography and recrystallized from EtOAc. m.p. 258-259° C.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  0.96 (t, 3,  $\text{CH}_3$ ,  $J=7$  Hz); 1.17 [m, 12,  $\text{O}-\text{CH}_2-\text{CH}_2-(\text{CH}_2)_6$ ]; 1.42 [m, 2,  $\text{O}-\text{CH}_2\text{CH}_2(\text{CH}_2)_6$ ]; 3.27-3.61  
20 (m, 4,  $\text{H}-5'$ ,  $\text{O}-\text{CH}_2(\text{CH}_2)_7$ ]; 3.95 (m, 1,  $\text{H}-4'$ ), 4.10-4.13 (m, 2,  $\text{H}-2'$ ,  $\text{H}-3'$ ); 5.13-6.06 (m, 2,  $3'-\text{OH}$   $5'-\text{OH}$ ); 5.80 (d, 1,  $\text{H}-1'$ ,  $J=6.4$  Hz); 6.47 (s, 2,  $2-\text{NH}_2$ ); 7.98 (s, 1,  $8-\text{H}$ ) and 10.64 (s, 1,  $\text{N}_1$  amide). Anal. Calcd. for  $\text{C}_{19}\text{H}_{31}\text{N}_5\text{O}_5$ : C, 55.73; H, 7.63; N, 17.10. Found: C, 55.67; H, 7.66; N, 17.02.

**25 EXAMPLE 14****N2-Isobutyryl-2'-O-nonylguanosine**

2'-O-nonylguanosine (14.7 g) in pyridine (360 ml) was treated with trimethylsilyl chloride (23.4 ml) and isobutyryl chloride (30.6 ml) as per the procedure of Example 4 to yield  
30 crude product (37 g). The crude material was purified by silica gel chromatography (eluted with 90/10  $\text{CHCl}_3/\text{MeOH}$ ) to yield 14.6 g of product re-crystallized from EtOAc. m.p. 168-169° C.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  0.85 [t, 3,  $\text{CH}_3$  (nonyl)], 1.14 [m, 18,  $\text{O}-\text{CH}_2\text{CH}_2(\text{CH}_2)_6$ ,  $\text{CH}(\text{CH}_3)_2$ ], 1.40 [m, 2,  $\text{O}-\text{CH}_2\text{CH}_2(\text{CH}_2)_6$ ], 2.79  
35 [m, 1,  $\text{CH}(\text{CH}_3)_2$ ], 3.31-3.63 (m, 4,  $\text{H}-5'$ ,  $\text{O}-\text{CH}_2(\text{CH}_2)_7$ ]; 3.96 (m,

- 33 -

1,  $\underline{\text{H}}\text{-4}'$ ), 4.27-4.37 (m, 2,  $\underline{\text{H}}\text{-2}'$ ,  $\underline{\text{H}}\text{-3}'$ ); 5.10 (t, 1,  $\underline{\text{H}}\text{-5}'$ ,  $\underline{\text{H}}\text{-OH}$ ,  $J=5$  Hz), 5.18 (d, 1,  $\underline{\text{H}}\text{-3}'$ ,  $\underline{\text{H}}\text{-OH}$ ,  $J=4$  Hz), 5.91 (d, 1,  $\underline{\text{H}}\text{-1}'$ ,  $J=6.6$  Hz), 8.31 (s, 1, 8- $\underline{\text{H}}$ ), 11.73 (s, 1,  $\text{C}_2$  amide) and 12.11 (s, 1,  $\text{N}_1$  amide). Anal. Calcd. for  $\text{C}_{23}\text{H}_{37}\text{N}_5\text{O}_6$ : C, 57.60; H, 7.78; N, 14.60. Found: C, 57.63; H, 7.92; N, 14.62.

**EXAMPLE 15****N2-Isobutyryl-5'-dimethoxytrityl-2'-O-nonylguanosine**

N2-Isobutyryl-2'-O-nonylguanosine (14.6 g, 30.4 mmol) was treated with dimethoxytrityl chloride (12.1 g, 34 mmol) in  
 10 pyridine (200 ml) as per the procedure of Example 5 to yield 16 g of purple foam prior to chromatography and 11.5 g after chromatography purification.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  0.84 [t, 3,  $\text{CH}_3$ (nonyl),  $J=7$  Hz], 1.16 [m, 18,  $\text{O}-\text{CH}_2\text{CH}_2(\text{CH}_2)_6$ ,  $\text{CH}(\text{CH}_3)_2$ ], 1.43 [m, 2,  $\text{O}-\text{CH}_2\text{CH}_2(\text{CH}_2)_6$ ], 2.77 [m, 1,  $\text{CH}(\text{CH}_3)_2$ ], 3.18-3.63 (m, 4,  
 15  $\underline{\text{H}}\text{-5}'$ ,  $\text{O}-\text{CH}_2(\text{CH}_2)_7$ ]; 3.74 (s, 6, DMTr  $\text{O}-\text{CH}_3$ ) 4.06 (m, 1,  $\underline{\text{H}}\text{-4}'$ ), 4.27 (m, 1,  $\underline{\text{H}}\text{-3}'$ ); 4.42 (m, 1,  $\underline{\text{H}}\text{-2}'$ ); 5.19 (d, 1,  $\underline{\text{H}}\text{-3}'$ ,  $\underline{\text{H}}\text{-OH}$ ,  $J=5$  Hz), 5.94 (d, 1,  $\underline{\text{H}}\text{-1}'$ ,  $J=5.7$  Hz), 6.83-7.38 (m, 13, DMTr aromatic), 8.14 (s, 1, 8- $\underline{\text{H}}$ ), 11.65 (s, 1,  $\text{C}_2$  amide) and 12.11 (s, 1,  $\text{N}_1$  amide). Anal. Calcd. for  $\text{C}_{44}\text{H}_{55}\text{N}_5\text{O}_8$ : C, 67.59; H, 7.27;  
 20 N, 8.96. Found: C, 67.59; H, 7.11; N, 8.80.

**EXAMPLE 16****N2-Isobutyryl-5'-dimethoxytrityl-2'-O-nonylguanosine 3'- $\beta$ -cyanoethyl-N,N-diisopropylphosphoramidite**

N2-Isobutyryl-5'-dimethoxytrityl-2'-O-nonylguanosine  
 25 (2.1 g) was treated with bis-(N,N-diisopropylamino)-2-cyanoethyl-phosphite (1.5 g) and N,N-diisopropylammonium tetrazolide (0.2 g) as per the procedure of Example 6 to yield the product (2.0 g).  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ )  $\delta$  150.7 and 150.4 (diastereomers).

30

**EXAMPLE 17****2,6-Diamino-9-(2',3'-di-O-propyl- $\beta$ -D-ribofuranosyl)purine**

The procedure of Example 2 was repeated utilizing 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine (10 g), NaH (3 g) and 1-  
 35 bromopropane (10 ml) in DMF. After evaporation of the reaction

- 34 -

solvent, the reaction products were purified by silica gel chromatography. The slower moving component yielded 4.3 g of the 2'-O-propyl product as a foam. This foam was crystallized from water to yield 3.6 g of product. The faster moving  
5 component isolated as an oil formed crystals upon standing. EtOH was added to the crystals, they were filtered and wash 1 x EtOH to yield 1.1 grams of 2',3'-di-O-propyl product. m.p. 165-167° C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.80 and 0.92 (t, 6, CH<sub>3</sub>), 1.6 and 1.45 (m, 4, CH<sub>2</sub>), 3.7-3.45 (br m, 6), 4.07 (m, 2), 4.5 (dd,  
10 1), 5.55 (br t, 1, 5'-OH), 5.8 (br s, 2, 6-NH<sub>2</sub>), 5.85 (d, 1, H-1'), 6.84 (br s, 2, 2-NH<sub>2</sub>) and 8.0 (s, 1, H-8).  
Anal. Calcd. for C<sub>16</sub>H<sub>26</sub>N<sub>6</sub>O<sub>4</sub>: C, 52.45; H, 7.15; N, 22.94. Found: C, 52.18; H, 7.19; N, 22.75.

**EXAMPLE 18****15 N2,N6-Diisobutyryl-2,6-diamino-9-(2'-O-propyl-β-D-ribofuranosyl)purine**

2,6-diamino-9-(2'-O-propyl-β-D-ribofuranosyl)purine (2.0 g) in pyridine (35 ml) was treated with trimethylsilyl chloride (3.9 ml, 5 eq) and isobutyryl chloride (3.2 ml, 5 eq)  
20 as per the procedure of Example 4 to yield a foam after silica gel chromatography. The foam was crystallized from EtOAc/Hex to yield 2.2 g of product. m.p. 140-142° C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.77 (t, 3, CH<sub>3</sub>), 1.07, 1.16 [d, 12, 2 x CH(CH<sub>3</sub>)<sub>2</sub>], 1.5 (m, 2, CH<sub>2</sub>), 2.9, 3.03 [m, 2, 2 x CH(CH<sub>3</sub>)<sub>2</sub>], 3.4 (m, 1, H-5''), 3.58  
25 (m, 3, OCH<sub>2</sub>, H-5'), 3.95 (m, 1, H-4'), 4.3 (m, 1), 4.5 (m, 1), 5.02 (t, 1, 5'-OH), 5.2 (d, 1, 3'-OH), 6.03 (d, 1, H-1'), 8.58 (s, 1, H-8), 10.39 (br s, 1, NH), and 10.57 (br s, 1, NH).

**EXAMPLE 19****30 N2,N6-Diisobutyryl-2,6-diamino-9-(5'-O-dimethoxytrityl-2'-O-propyl-β-D-ribofuranosyl)purine**

N2,N6-Diisobutyryl-2,6-diamino-9-(2'-O-propyl-β-D-ribo-furanosyl)purine (1.9 g) was treated with dimethoxytrityl chloride (1.5 g, 1.1 eq), and dimethylaminopyridine (20 mg as a catalyst) in pyridine (50 ml) as per the procedure of Example  
35 5 to yield the product as a foam (2.8 g). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ

- 35 -

0.79 (t, 3,  $\text{CH}_3$ ), 1.07, 1.16 [d, 12, 2 x  $\text{CH}(\text{CH}_3)_2$ ], 1.5 (m, 2,  $\text{CH}_2$ ), 2.9, 3.03 [m, 2, 2 x  $\text{CH}(\text{CH}_3)_2$ ], 3.58 (m, 3,  $\text{OCH}_2$ ,  $\text{H}-5'$ ), 4.15 (m, 1,  $\text{H}-4'$ ), 4.4 (m, 1), 4.6 (m, 1), 5.15 (d, 1,  $3'-\text{OH}$ ), 6.15 (d, 1,  $\text{H}-1'$ ), 6.—

5 8-7.35 (m, 13, DMTr), 8.5 (s, 1,  $\text{H}-8$ ), 10.3 (br s, 1,  $\text{NH}$ ), and 10.57 (br s, 1,  $\text{NH}$ ).

**EXAMPLE 20**

**N2,N6-Diisobutyryl-2,6-diamino-9-(5'-O-dimethoxytrityl-2'-O-propyl- $\beta$ -D-ribofuranosyl)purine 3'- $\beta$ -cyanoethyl-N,N-**  
 10 **diisopropylphosphoramidite**

N2,N6-Diisobutyryl-2,6-diamino-9-(5'-O-dimethoxytrityl-2'-O-propyl- $\beta$ -D-ribofuranosyl)purine (2.6 g) was treated with bis-(N,N-diisopropylamino)-2-cyanoethylphosphite (1.7 g) and N,N-diisopropylammonium tetrazolide (300 mg) overnight at  
 15 room temperature. The reaction mixture was partitioned against dil.  $\text{Na}_2\text{CO}_3/\text{CHCl}_2$  and then  $\text{Na}_2\text{CO}_3/\text{NaCl}$  and dried over  $\text{MgSO}_4$ . The organic layer was evaporated to a foam. The foam was dissolved in  $\text{CH}_2\text{Cl}_2$  ( $\approx 8$  ml) and slowly added to Hexanes (500 ml). The solid was filtered and dried to yield the product as a powder  
 20 (3.1 g).  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ )  $\delta$  150.8 and 151.3.

**EXAMPLE 21**

**2,6-Diamino-9-[2'-O-[(N-phthalimido)prop-3-yl]- $\beta$ -D-ribofuranosyl] purine & 2,6-Diamino-9-[3'-O-[(N-phthalimido)prop-3-yl]- $\beta$ -D-ribo-furanosyl]purine**  
 25 2,6-Diamino-9-( $\beta$ -D-ribofuranosyl)purine (14.2 g) was treated with sodium hydride (3 g, 1.5 eq) and N-(3-bromopropyl)phthalimide (5.3 ml, 1.5 eq) in DMF (20 g) at 70° C overnight. The reaction mixture was proportioned between  $\text{H}_2\text{O}$  and Hexanes (1x) and the aqueous layer then extracted 4 x  $\text{CH}_2\text{Cl}_2$ . The  
 30 organic layer was dried over  $\text{MgSO}_4$  and evaporated to a residue. The residue was purified by silica gel chromatography eluted with  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ . The 2'-O-(N-phthalimido)propyl product eluted first followed by mixed fractions and then the 3'-O-(N-phthalimido) product. Evaporations of the fractions gave 3.4  
 35 g of the 2'-O-(N-phthalimido)propyl product, 3.0 g of mixed 2'

- 36 -

and 3' products and 1.4 g of the 3'-O-(N-phthalimido)propyl product all as foams. The 3'-O-(N-phthalimido)propyl product was crystallized from EtOAc/MeOH to give 270 mg of solid.

**2,6-Diamino-9-[2'-O-[(N-phthalimido)prop-3-yl]- $\beta$ -D-ribofuranosyl] purine**

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  1.8 (tq, 2, -CH<sub>2</sub>-), 3.4-3.58 (m, 6, 2x CH<sub>2</sub>, H-5'), 3.9 (m, 1), 4.26 (m, 1), 4.37 (m, 1), 5.05 (br d, 1, 3'-OH), 5.4 (br t, 1, 5'-OH), 5.72 (br s, 2, NH<sub>2</sub>), 5.8 (br d, 1, H-1'), 6.75 (br s, 2, NH<sub>2</sub>), 7.8 (br s, 4, Ar) and 8.93 (s, 1, H-8).

**2,6-Diamino-9-[3'-O-[(N-phthalimido)prop-3-yl]- $\beta$ -D-ribofuranosyl] purine**

m.p. 220-222° C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  1.85 (tq, 2, -CH<sub>2</sub>-N), 3.6-3.67 (m, 4, -O-CH<sub>2</sub>, H-5'), 3.85 (m, 1), 3.92 (m, 1), 4.6 (m, 1), 5.33 (d, 1, 2'-OH), 5.45 (br t, 1, 5'-OH), 5.65 (d, 1, H-1'), 5.73 (br s, 2, NH<sub>2</sub>), 6.75 (br d, 2, NH<sub>2</sub>), 7.8-7.85 (m, 4, Ar) and 7.85 (s, 1, H-8). Anal. Calcd. for C<sub>21</sub>H<sub>23</sub>N<sub>7</sub>O<sub>6</sub>: C, 53.73; H, 4.94; N, 20.88. Found: C, 53.59; H, 4.89; N, 20.63.

**EXAMPLE 22**

**2'-O-[(N-Phthalimido)prop-3-yl] guanosine**

2,6-diamino-9-[2'-O-[(N-phthalimido)prop-3-yl]- $\beta$ -D-ribofuranosyl] purine (3.1 g) in 0.1 M sodium phosphate buffer (3 ml, pH 7.4), 0.05 M tris buffer (65 ml, pH 7.4) and DMSO (45 ml) was treated with adenosine deaminase (200 mg) at room temperature for 5 days as per the procedure of Example 3. The product containing fractions from the silica gel chromatography were evaporated and upon concentration formed white crystals. The crystals were filtered and washed with MeOH to yield 1.1 g of product. An analytical sample was recrystallized from MeOH.

m.p. 192-194° C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  1.82 (m, 2, CH<sub>2</sub>), 3.45-3.67 (m, 6, H-5', OCH<sub>2</sub>, NCH<sub>2</sub>), 3.9 (m, 1), 4.3 (m, 2, H-2', H-3'), 5.1 (m, 2, 5' and 3'-OH), 5.8 (d, 1, H-1'), 6.5 (br s, 2, NH<sub>2</sub>), 7.83 (s, 4, phthal), 7.98 (s, 1, H-8) and 10.5 (br s, 1, NH). Anal. Calcd. for C<sub>21</sub>H<sub>22</sub>N<sub>6</sub>O<sub>7</sub>·½H<sub>2</sub>O: C, 52.61; H, 4.83; N, 17.53. Found: C, 52.52; H, 4.78; N, 17.38.

**EXAMPLE 23****N2-Isobutyryl-2'-O-[(N-phthalimido)prop-3-yl] guanosine**

2'-O-[(N-phthalimido)prop-3-yl] guanosine (7.2 g, crude) in pyridine (35 ml) was treated with trimethylsilyl chloride (11.6 ml, 5 eq) and isobutyryl chloride (8 ml, 5 eq) as per the procedure of Example 4 to yield the product as a crude foam (6.5 g). An analytical sample was obtained by crystallization from EtOAc. m.p. 166-168° C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.15 [d, 6, -CH(CH<sub>3</sub>)<sub>2</sub>], 1.85 (m, 2, CH<sub>2</sub>), 2.8 [m, 1, CH(CH<sub>3</sub>)<sub>2</sub>], 3.45-3.7 (m, 6, H-5', OCH<sub>2</sub>, NCH<sub>2</sub>), 3.95 (m, 1), 4.34 (m, 1), 4.4 (m, 1), 5.12 (t, 1, 5'-OH), 5.18 (d, 1, 3'-OH), 5.9 (d, 1, H-1'), 7.83 (s, 4, phthal), 8.3 (s, 1, H-8), 11.65 (br s, 1, NH) and 12.1 (br s, 1, NH). Anal. Calcd. for C<sub>25</sub>H<sub>28</sub>N<sub>6</sub>O<sub>8</sub>·½H<sub>2</sub>O: C, 54.64; H, 5.32; N, 15.29. Found: C, 54.46; H, 5.39; N, 14.98.

15

**EXAMPLE 24****N2-Isobutyryl-5'-dimethoxytrityl-2'-O-[(N-phthalimido)prop-3-yl]guanosine**

N2-Isobutyryl-2'-O-[(N-phthalimido)prop-3-yl] guanosine (1.2 g) was treated with dimethoxytrityl chloride (820 mg, 1.1 eq), and dimethylaminopyridine (20 mg as a catalyst) in pyridine (50 ml) as per the procedure of Example 5 utilizing 1:1 Hex/EtOAc, then EtOAc then 5%MeOH/EtOAc with 1% TEA as eluent. The product containing fraction were evaporated to yield the product as a foam (1.7 g). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.1 [d, 6, -CH(CH<sub>3</sub>)<sub>2</sub>], 1.85 (m, 2, CH<sub>2</sub>), 2.75 [m, 1, CH(CH<sub>3</sub>)<sub>2</sub>], 3.45-3.7 (m, 6, H-5', OCH<sub>2</sub>, NCH<sub>2</sub>), 3.75 (s, 6, OCH<sub>3</sub>), 4.0 (m, 1), 4.32 (m, 1), 4.4 (m, 1), 5.2 (d, 1, 3'-OH), 5.93 (d, 1, H-1'), 6.83, 7.2, 7.35 (m, 13, DMTr), 7.78 (s, 4, phthal), 8.15 (s, 1, H-8), 11.6 (br s, 1, NH) and 12.05 (br s, 1, NH). Anal. Calcd. for C<sub>46</sub>H<sub>46</sub>N<sub>6</sub>O<sub>10</sub>·H<sub>2</sub>O: C, 64.18; H, 5.62; N, 9.76. Found: C, 64.42; H, 5.78; N, 9.53.

25  
30

- 38 -

**EXAMPLE 25**

**N2-Isobutyryl-5'-dimethoxytrityl-2'-O-[(N-phthalimido)prop-3-yl]guanosine 3'-β-cyanoethyl-N,N-diisopropylphosphoramidite**

N2-Isobutyryl-5'-dimethoxytrityl-2'-O-[(N-phthalimido)prop-3-yl] guanosine (1.6 g) was treated with *bis*-(N,N-diisopropylamino)-2-cyanoethylphosphite (1.48 g) and N,N-diisopropylammonium tetrazolide (200 mg) as per the procedure of Example 6 to yield the product (2.0 g). <sup>31</sup>P NMR (CDCl<sub>3</sub>) δ 150.9.

**10 EXAMPLE 26**

**N2-Dimethylaminomethylidene-5'-dimethoxytrityl-2'-O-[(N-phthalimido)prop-3-yl] guanosine**

2'-O-[(N-phthalimido)prop-3-yl] guanosine (900 mg) in DMF (20 ml) was treated with N,N-dimethylformamide dimethyl acetal (2 ml). The reaction mixture was stirred for 2 hr and evaporated under high vac at 52° C. The residue was co-evaporated 1 x with pyridine and taken up in solution in pyridine. Dimethoxytrityl chloride (713 mg, 1.1 eq) and dimethylaminopyridine (20 mg as a catalyst) were added. The reaction mixture was stirred overnight, partitioned between Na<sub>2</sub>CO<sub>3</sub>/CH<sub>2</sub>Cl<sub>2</sub>, dried over MgSO<sub>4</sub> and purified by silica gel chromatography as per the procedure of Example 5 to yield 1.7 g of product as an off white solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.88 (m, 2, CH<sub>2</sub>), 3.1 [d, 6, N=CHN(CH<sub>3</sub>)<sub>2</sub>], 3.3 (m, 2, H-5'), 3.67 (m, 4, OCH<sub>2</sub>, NC<sub>2</sub>), 3.78 (s, 6, 2x OCH<sub>3</sub>), 4.0 (m, 1, H-4'), 4.35 (m, 2, H-2', H-3'), 5.2 (d, 1, 3'-OH), 5.95 (d, 1, H-1'), 6.85, 7.25, 7.39 (m, 13, DMTr), 7.85 (s, 4, phthal), 7.95 [s, 1, H-8), 8.5 (s, 1, N=CHN(CH<sub>3</sub>)<sub>2</sub>] and 11.39 (s, 1, NH<sub>2</sub>). Anal. Calcd. for C<sub>45</sub>H<sub>45</sub>N<sub>7</sub>O<sub>9</sub>·½H<sub>2</sub>O: C, 64.58; H, 5.54; N, 11.71. Found: C, 64.10; H, 5.65; N, 11.47.



**EXAMPLE 27**

**N2-Dimethylaminomethylidene-5'-dimethoxytrityl-2'-O-[(N-phthalimido)prop-3-yl] guanosine 3'-β-cyanoethyl-N,N-diisopropylphosphoramidite**

5           N2-dimethylaminomethylidene-5'-dimethoxytrityl-2'-O-[(N-phthalimido)prop-3-yl] guanosine (1.7 g), bis-(N,N-diisopropylamino)-2-cyanoethylphosphite (1.4 ml) and N,N-diisopropylammonium tetrazolide (170 mg) were stirred overnight at room temperature. The reaction mixture was partitioned  
10 between CH<sub>2</sub>Cl<sub>2</sub> and Na<sub>2</sub>CO<sub>3</sub> (2 x). The organic phase was dried over MgSO<sub>4</sub> and evaporated to an oil. The oil was dissolved in a minimum of CH<sub>2</sub>Cl<sub>2</sub> and added dropwise to ≈ 900 ml Hexanes to precipitate the product. The solid was isolated and dried to yield 2.1 g of product. <sup>1</sup>P NMR (CDCl<sub>3</sub>) δ 150.4, 150.6.

**15 EXAMPLE 28**

**2,6-Diamino-9-[2'-O-(N-phthalimido)pent-5-yl]-β-D-ribofuranosyl purine**

2,6-Diamino-(9-β-D-ribofuranosyl)purine (6.7 g) was treated with sodium hydride (1.3 g) and N-(5-bromopentyl)  
20 phthalimide (7.8 g, 1.1 eq) in DMF (60 ml) at room temperature for three days. The reaction mixture was proportioned between H<sub>2</sub>O and CH<sub>2</sub>Cl<sub>2</sub> and extracted 4 x CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layers were dried over MgSO<sub>4</sub> and evaporated. The residue was purified by silica gel chromatography eluted with 5 → 10%  
25 MeOH/CH<sub>2</sub>Cl<sub>2</sub>. The 2'-O-(N-phthalimido)pentyl containing fractions were collected and evaporated to a yellow foam to give 2.2 g of product. An analytical sample was crystallized from EtOH. m.p. 173-175° C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.2 (m, 2, -CH<sub>2</sub>-), 1.47 (m, 4, 2x CH<sub>2</sub>), 3.55, 3.65 (m, 6, O-CH<sub>2</sub>, H-5',  
30 NCH<sub>2</sub>), 3.95 (m, 1), 4.28 (m, 1), 4.4 (m, 1), 5.13 (d, 1, 3'-OH), 5.5 (t, 1, 5'-OH), 5.77 (br s, 2, 6-NH<sub>2</sub>), 5.84 (br d, 1, H-1'), 6.8 (br s, 2, 2-NH<sub>2</sub>), 7.86 (M, 4, phthal) and 7.95 (s, 1, H-8). Anal. Calcd. for C<sub>23</sub>H<sub>27</sub>N<sub>7</sub>O<sub>6</sub>: C, 55.50; H, 5.47; N, 19.71. Found: C, 55.44; H, 5.51; N, 19.30.

**EXAMPLE 29****2'-O-[(N-Phthalimido)pent-5-yl] guanosine**

A mixture of the 2,6-diamino-9-[2'-O-[(N-phthalimido)pent-5-yl]- $\beta$ -D-ribofuranosyl]purine and 2,6-diamino-9-[3'-O-[(N-phthalimido)pent-5-yl]- $\beta$ -D-ribofuranosyl]purine isomers (2.2 g) in 0.1 M tris buffer (60 ml, pH 7.4), 0.1 M NaPO<sub>4</sub> buffer (2 ml, pH 7.4) and DMSO (40 ml) was treated with adenosine deaminase (60 mg) at room temperature for 5 days as per the procedure of Example 3. The product containing fractions from the silica gel chromatography were evaporated to give the product (1.0 g) as a crude white solid. An analytical sample was prepared by the addition of MeOH to form crystals. m.p. 178-180° C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  1.24 (m, 2, CH<sub>2</sub>), 1.5 (m, 4, 2x CH<sub>2</sub>), 3.5-3.6 (m, 6, H-5', OCH<sub>2</sub>, NCH<sub>2</sub>), 3.87 (m, 1, H-4'), 4.25 (m, 2, H-2', H-3'), 5.1 (m, 2, 5' and 3'-OH), 5.78 (d, 1, H-1'), 6.5 (br s, 2, NH<sub>2</sub>), 7.84 (M, 4, phthal), 7.98 (s, 1, H-8) and 10.67 (br s, 1, NH). Anal. Calcd. for C<sub>23</sub>H<sub>26</sub>N<sub>6</sub>O<sub>7</sub>·½H<sub>2</sub>O: C, 54.43; H, 5.36; N, 16.56. Found: C, 54.79; H, 5.24; N, 16.61.

**20 EXAMPLE 30****N2-Isobutyryl-2'-O-[(N-phthalimido)pent-5-yl] guanosine**

2'-O-[(N-phthalimido)pent-5-yl] guanosine (1.6 g, crude) in pyridine (35 ml) was treated with trimethylsilyl chloride (2.0 ml, 5 eq) and isobutyryl chloride (1.68 ml, 5 eq) as per the procedure of Example 4 to yield the product as a foam. This foam was co-evaporated 2 x with EtOAc followed by the addition of EtOAc and heating to yield white crystals (950 mg). m.p. 202-204° C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  1.1 [d, 6, -CH(CH<sub>3</sub>)<sub>2</sub>], 1.17 (m, 2, CH<sub>2</sub>), 1.43 (m, 4, 2x CH<sub>2</sub>), 2.74 [m, 1, CH(CH<sub>3</sub>)<sub>2</sub>], 3.45-3.55 (m, 6, H-5', OCH<sub>2</sub>, NCH<sub>2</sub>), 3.9 (m, 1), 4.25 (m, 1), 4.3 (m, 1), 5.07 (t, 1, 5'-OH), 5.15 (d, 1, 3'-OH), 5.87 (d, 1, H-1'), 7.8 (s, 4, phthal), 8.27 (s, 1, H-8), 11.67 (br s, 1, NH) and 12.06 (br s, 1, NH). Anal. Calcd. for C<sub>27</sub>H<sub>32</sub>N<sub>6</sub>O<sub>8</sub>·½H<sub>2</sub>O: C, 56.14; H, 5.76; N, 14.55. Found: C, 56.45; H, 5.74; N, 14.41.

- 41 -

**EXAMPLE 31**

**N2-Isobutyryl-5'-dimethoxytrityl-2'-O-[(N-phthalimido)pent-5-yl] guanosine**

N2-Isobutyryl-2'-O-[(N-phthalimido)pent-5-yl]  
5 guanosine (0.95 g) was treated with dimethoxytrityl chloride (620 mg, 1.1 eq), and dimethylaminopyridine (20 mg as a catalyst) in pyridine (50 ml) as per the procedure of Example 5 utilizing EtOAc 1% TEA and then 5% MeOH EtOAc/CH<sub>2</sub>Cl<sub>2</sub> with 1% TEA as eluent. The product containing fractions were  
10 evaporated to yield the product as a foam (1.4 g). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.14 [d, 6, -CH(CH<sub>3</sub>)<sub>2</sub>], 1.25 (m, 2, CH<sub>2</sub>), 1.53 (m, 4, 2x CH<sub>2</sub>), 2.77 [m, 1, CH(CH<sub>3</sub>)<sub>2</sub>], 3.3-3.6 (m, 6, H-5', OCH<sub>2</sub>, NCH<sub>2</sub>), 3.75 (s, 6, OCH<sub>3</sub>), 4.07 (m, 1), 4.33 (m, 1), 4.4 (m, 1), 5.18 (d, 1, 3'-OH), 5.94 (d, 1, H-1'), 6.83, 7.2, 7.53 (m, 13,  
15 DMTr), 7.8 (s, 4, phthal), 8.15 (s, 1, H-8), 11.6 (br s, 1, NH) and 12.1 (br s, 1, NH). Anal. Calcd. for C<sub>48</sub>H<sub>50</sub>N<sub>6</sub>O<sub>10</sub>·½H<sub>2</sub>O: C, 65.52; H, 5.84; N, 9.55. Found: C, 65.55; H, 5.94; N, 9.20.

**EXAMPLE 32**

**2,6-Diamino-9-[3',5'-O-(tetraisopropylidisiloxane-1,3-diyl)-β-D-  
20 ribofuranosyl]purine**

To a suspension of 2,6-diamino-9-(β-D-ribofuranosyl)purine (10.5 g) in pyridine (100 ml) was added 1,3-dichlorotetraisopropylidisiloxane (TIPDS, 12.6 g). The reaction was stirred at room temperature for 4 hours and an additional  
25 1.3 g of 1,3-dichlorotetraisopropylidisiloxane was added followed by stirring overnight. The reaction mixture was poured into ice water and the insoluble product (11.6 g) collected by filtration. An analytical sample was recrystallized from EtOAc/Hexanes. m.p. 170-172° C. Anal.  
30 Calcd. for C<sub>22</sub>H<sub>40</sub>N<sub>6</sub>O<sub>5</sub>Si<sub>2</sub>·½H<sub>2</sub>O: C, 49.5; H, 7.74; N, 15.7. Found: 49.57; H, 7.82; N, 15.59.

- 42 -

**EXAMPLE 33****2,6-Diamino-9-[3',5'-O-(tetraisopropylidisiloxane-1,3-diyl)-2-O-methyl- $\beta$ -D-ribofuranosyl]purine**

A mixture of 2,6-Diamino-9-[3,5-O-(tetraisopropylidisiloxane-1,3-diyl)- $\beta$ -D-ribofuranosyl]purine (8.8 g) in DMF (120 ml) and methyl iodide (3 ml, 3 eq) was cooled in an ice bath and NaH (60% in oil, 1.0 g, 1.5 eq) added. After 20 min the reaction was quenched with MeOH and partitioned between sat.  $\text{NH}_4\text{Cl}$  and  $\text{CH}_2\text{Cl}_2$ . The organic phase was washed with 1 x  $\text{NH}_4\text{Cl}$ , dried over  $\text{MgSO}_4$  and evaporated. The residue was crystallized from hot EtOH/ $\text{H}_2\text{O}$  to yield the product (8.5 g) as crystals. m.p. 87-89°C.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  1.05 (m, 28, TIPDS), 3.57 (s, 3,  $\text{OCH}_3$ ), 3.98 (m, 1,  $\text{H}-4'$ ), 3.92 and 4.07 (ABX, 2,  $\text{H}-5'$ ), 4.13 (d, 1), 4.6 (dd, 1,  $\text{H}-3'$ ), 5.76 (br s, 2,  $\text{NH}_2$ ), 5.8 (s, 1,  $\text{H}-1'$ ), 6.77 (br s, 2,  $\text{NH}_2$ ) AND 7.77 (s, 1  $\text{H}-8$ ).

**EXAMPLE 34****2,6-Diamino-9-(2'-O-methyl- $\beta$ -D-ribofuranosyl)purine**

To a solution of 2,6-Diamino-9-[3',5'-O-(tetraisopropylidisiloxane-1,3-diyl)-2'-O-methyl- $\beta$ -D-ribofuranosyl]purine (8.5 g) in THF (50 ml) was added 1M tetrabutylammonium fluoride in THF (Aldrich, 20 ml). The reaction mixture was stirred for 2 hrs and filtered. The filter cake was washed with 2 x EtOAc and air dried to give 4.0 g of crude product. An analytical sample was crystallized from hot MeOH. m.p. 133-135°C.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  3.3 (s, 3,  $\text{OCH}_3$ ), 3.58 (m, 2,  $\text{H}-5'$ ), 3.98 (m, 1,  $\text{H}-4'$ ), 4.28 (m, 2,  $\text{H}-2'$ ,  $\text{H}-3'$ ), 5.23 (br s, 1, 3'-OH), 5.48 (br t, 1, 5'-OH), 5.77 (br s, 2,  $\text{NH}_2$ ), 5.82 (d, 1,  $\text{H}-1'$ ), 6.83 (br s, 2,  $\text{NH}_2$ ) and 7.95 (s, 1,  $\text{H}-8$ ). Anal. Calcd. for  $\text{C}_{11}\text{H}_{16}\text{N}_6\text{O}_4 \cdot \frac{1}{2}\text{H}_2\text{O}$ : C, 43.28; H, 5.61; N, 27.52. Found: C, 43.51; H, 5.62; N, 27.26.

**EXAMPLE 35****2'-O-Methylguanosine**

2,6-Diamino-9-(2'-O-methyl- $\beta$ -D-ribofuranosyl)purine (9.5 g) in 0.1M sodium phosphate buffer (200 ml, pH 7.4) and DMSO (25 ml) was treated with adenosine deaminase (Type II Sigma) at RT for 4 days. The resulting suspension was cooled and filtered and the resulting filter cake washed with H<sub>2</sub>O and dried to a white solid (4.0 g). The solid was recrystallized from hot H<sub>2</sub>O to yield 2.9 g of product. m.p. 236-238° C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  3.3 (s, 3, OCH<sub>3</sub>), 3.53 and 3.6 (ABX, 2, H-5'), 3.87 (m, 1, H-4'), 4.15 (m, 1, H-2'), 4.25 (m, 1, H-3'), 5.13 (t, 1, 5'-OH), 5.23 (d, 1, 3'-OH), 5.8 (d, 1, H-1'), 6.48 (br s, 2, NH<sub>2</sub>), 7.96 (s, 1, H-8) and 10.68 (br s, 1, NH). Anal. Calcd. for C<sub>11</sub>H<sub>15</sub>N<sub>5</sub>O<sub>5</sub>·½H<sub>2</sub>O: C, 43.14; H, 5.26; N, 22.86. Found: C, 43.59; H, 5.34; N, 23.04.

**EXAMPLE 36****N2-Isobutyryl-2'-O-methylguanosine**

2'-O-methylguanosine (3.5 g) in pyridine (100 ml) was treated with trimethylsilyl chloride (9 ml, 6 eq) and isobutyryl chloride (6.2 ml) at RT for 4 hr. The reaction mixture was cooled in an ice bath, H<sub>2</sub>O (20 ml) was added and stirring continued for an additional 20 min. NH<sub>4</sub>OH (20 ml) was added and after stirring for 30 min the reaction mixture was evaporated. The residue was triturated with H<sub>2</sub>O, filtered and the filtrate evaporated and purified by silica gel chromatography as per the procedure of Example 4 to yield the product as an off white solid (1.5 g). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  1.1 [d, 6, CH(CH<sub>3</sub>)<sub>2</sub>], 2.77 [m, 1, CH(CH<sub>3</sub>)<sub>2</sub>], 3.33-3.6 (m, 5, OCH<sub>3</sub>, H-5'), 3.93 (m, 1, H-4'), 4.22 (m, 1), 4.3 (m, 1), 5.1 (t, 1, 5'-OH), 5.28 (d, 1, 3'-OH), 5.9 (d, 1, H-1'), 8.28 (s, 1, H-8) and 11.9 (br s, 1, NH).

- 44 -

**EXAMPLE 37****N2-Isobutyryl-5'-dimethoxytrityl-2'-O-methylguanosine**

N2-Isobutyryl-2'-O-methylguanosine (1.5 g) was treated with dimethoxytrityl chloride (1.5 g, 1.1 eq), and  
5 dimethylaminopyridine (100 mg as a catalyst) in pyridine (50 ml) as per the procedure of Example 5 to yield the product as a foam (2.6 g). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.14 (d, 6, CH(CH<sub>3</sub>)<sub>2</sub>), 2.75 [m, 1, CH(CH<sub>3</sub>)<sub>2</sub>], 3.5 (m, 2, H-5'), 3.74 (s, 6, OCH<sub>3</sub>), 4.05 (m, 1), 4.33 (m, 1), 5.26 (d, 1, 3'-OH), 5.95 (d, 1, H-1'),  
10 7.2, 7.35 (m, 13, DMTr), 8.15 (s, 1, H-8), 11.6 (br s, 1, NH) and 12.1 (br s, 1, NH).

**EXAMPLE 38****N2-Isobutyryl-5'-dimethoxytrityl-2'-O-methylguanosine 3'-β-cyanoethyl-N,N-diisopropylphosphoramidite**

15 N2-Isobutyryl-5'-dimethoxytrityl-2'-O-methylguanosine (20 g) was treated with bis-(N,N-diisopropylamino)-2-cyanoethylphosphite (10.8 g) and N,N-diisopropylammonium tetrazolide (1.6 g) as per the procedure of Example 6 to yield the product (15.7 g). <sup>31</sup>P NMR (CDCl<sub>3</sub>) δ 148.97 and 147.96.

**20 EXAMPLE 39****N2,N6-Diisobutyryl-2,6-diamino-9-(2'-O-methyl-β-D-ribofuranosyl) purine**

2,6-diamino-9-(2'-O-methyl-β-D-ribofuranosyl)purine (700 mg) in pyridine (20 ml) was treated with trimethylsilyl  
25 chloride (2.1 ml, 7 eq) and isobutyryl chloride (1.25 ml, 5 eq) as per the procedure of Example 4 to yield the product as a foam (900 mg) after silica gel chromatography.

**EXAMPLE 40****30 N2,N6-Diisobutyryl-2,6-diamino-9-(5'-O-dimethoxytrityl-2'-O-methyl-β-D-ribofuranosyl)purine**

N2,N6-Diisobutyryl-2,6-diamino-9-(2'-O-methyl-β-D-ribofuranosyl)purine (900 mg) was treated with dimethoxytrityl chloride (1.0 g) and dimethylaminopyridine (20 mg as a catalyst) in pyridine (30 m) as per the procedure of Example 5

- 45 -

to yield the product as a foam (700 mg).  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  0.96-1.16 [m, 12, 2x  $\text{CH}(\text{CH}_3)_2$ ], 2.9 and 3.05 [M, 2, 2x  $\text{CH}(\text{CH}_3)_2$ ], 3.18 and 3.37 (ABX, 2,  $\text{H}-5'$ ), 3.38 (s, 3,  $\text{OCH}_3$ ), 3.7 (s, 6,  $\text{OCH}_3$ ), 4.05 (m, 1,  $\text{H}-4'$ ), 4.44 (m, 2,  $\text{H}-2', \text{H}-3'$ ), 5.24 (d, 1,  $3'-\text{OH}$ ), 6.06 (d, 1,  $\text{H}-1'$ ), 6.78, 7.2, 7.33 (m, 13, Ar), 8.22 (s, 1,  $\text{H}-8$ ), 10.3 (br s, 1,  $\text{NH}$ ) and 10.57 (br s, 1,  $\text{NH}$ ).

**EXAMPLE 41**

**N2,N6-Diisobutyryl-2,6-diamino-9-(5'-O-dimethoxytrityl-2'-O-methyl- $\beta$ -D-ribofuranosyl)purine 3'- $\beta$ -cyanoethyl-N,N-diisopropylphosphoramidite**

N2,N6-Diisobutyryl-2,6-diamino-9-(5'-O-dimethoxytrityl-2'-O-methyl- $\beta$ -D-ribofuranosyl)purine (600 mg) was treated with *bis*-(N,N-diisopropylamino)-2-cyanoethylphosphite (500  $\mu\text{l}$ ) and N,N-diisopropylammonium tetrazolide (80 mg) overnight at RT. The reaction mixture was partitioned against dil.  $\text{Na}_2\text{CO}_3/\text{CHCl}_2$  and then  $\text{Na}_2\text{CO}_3/\text{NaCl}$  and dried over  $\text{MgSO}_4$ . The organic layer was evaporated to a foam (500 mg).  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ )  $\delta$  151.1 (doublet).

**EXAMPLE 42**

**2,6-Diamino-9-(2'-O-octadecyl- $\beta$ -D-ribofuranosyl)purine**

2,6-Diamino-9-( $\beta$ -D-ribofuranosyl)purine (50 g, 180 mmol) and sodium hydride (7 g) in DMF (1 l) were heated to boiling for 2 hr. Iodoctadecane (100 g) was added at  $150^\circ\text{C}$  and the reaction mixture allowed to cool to RT. The reaction mixture was stirred for 11 days at RT. The solvent was evaporated and the residue purified by silica gel chromatography. The product was eluted with 5%  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ . The product containing fraction were evaporated to yield the product (11 g).  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  0.84 (t, 3,  $\text{CH}_2$ ); 1.22 [m, 32,  $\text{O}-\text{CH}_2-\text{CH}_2-(\text{CH}_2)_{16}-$ ]; 1.86 (m, 2,  $\text{O}-\text{CH}_2\text{CH}_2-$ ); 3.25 (m, 2,  $\text{O}-\text{CH}_2-$ ); 3.93 (d, 1,  $4'\text{H}$ ), 4.25 (m, 1,  $3'\text{H}$ ); 4.38 (t, 1,  $2'\text{H}$ ); 5.08 (d, 1,  $3'-\text{OH}$ ); 5.48 (t, 1,  $5'-\text{OH}$ ); 5.75 (s, 2,  $6-\text{NH}_2$ ); 5.84 (d, 1,  $1'-\text{H}$ ); 6.8 (s, 2,  $2-\text{NH}_2$ ); and 7.95 (s, 1,  $8-\text{H}$ ).

**EXAMPLE 43****2'-O-Octadecylguanosine**

2,6-Diamino-9-(2'-O-octadecyl- $\beta$ -D-ribofuranosyl) purine (10 g) in 0.1 M sodium phosphate buffer (50 ml, pH 7.4),  
5 0.1 M tris buffer (1000 ml, pH 7.4) and DMSO (1000 ml) was treated with adenosine deaminase (1.5 g) as per the procedure of Example 3. At day 3, day 5 and day 7 an additional aliquot (500 mg, 880 mg and 200 mg, respectively) of adenosine deaminase was added. The reaction was stirred for a total of  
10 9 day and after purification by silica gel chromatography yielded the product (2 g). An analytical sample was recrystallized from MeOH  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  0.84 (t, 3,  $\text{CH}_3$ ), 1.22 [s, 32, O- $\text{CH}_2$ - $\text{CH}_2$ -( $\text{CH}_2$ )<sub>16</sub>], 5.07 (m, 2, 3'-OH 5'-OH); 5.78 (d, 1, 1'-H); 6.43 (s, 2,  $\text{NH}_2$ ), 7.97 (s, 1, 8-H) and 10.64 (s, 1,  $\text{NH}_2$ ).  
15 Anal. Calcd. for  $\text{C}_{28}\text{H}_{49}\text{N}_5\text{O}_5$ : C, 62.80; H, 9.16; N, 12.95. Found: C, 62.54; H, 9.18; N, 12.95.

**EXAMPLE 44****N2-Isobutyryl-2'-O-octadecylguanosine**

2'-O-Octadecylguanosine (1.9 g) in pyridine (150 ml)  
20 was treated with trimethylsilyl chloride (2 g, 5 eq) and isobutyryl chloride (2 g, 5 eq) as per the procedure of Example 4. The product was purified by silica gel chromatography (eluted with 3% MeOH/EtOAc) to yield 1.2 g of product.  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  0.85 [t, 3,  $\text{CH}_3$ ], 1.15 [m, 38, O- $\text{CH}_2$  $\text{CH}_2$ -( $\text{CH}_2$ )<sub>16</sub>,  
25 CH( $\text{CH}_3$ )<sub>2</sub>], 2.77 [m, 1, CH( $\text{CH}_3$ )<sub>2</sub>], 4.25 (m, 2, 2'-H, 3'-H); 5.08 (t, 1, 5'-OH), 5.12 (d, 1, 3'-OH), 5.87 (d, 1, 1'-H), 8.27 (s, 1, 8-H), 11.68 (s, 1,  $\text{NH}_2$ ) and 12.08 (s, 1,  $\text{NH}_2$ ). Anal. Calcd. for  $\text{C}_{32}\text{H}_{55}\text{N}_5\text{O}_6$ : C, 63.47; H, 9.09; N, 11.57. Found: C, 63.53; H, 9.20; N, 11.52.

**30 EXAMPLE 45****2,6-Diamino-9-[2'-O-(imidazol-1-yl)butyl- $\beta$ -D-ribofuranosyl] purine**

2,6-Diamino-(9- $\beta$ -D-ribofuranosyl)purine (5.0 g) in DMF (400 ml) was treated with sodium hydride (0.78 g). After  
35 stirring an additional 30 min a further portion of sodium



- 47 -

hydride (2.6 g) was added immediately followed by bromobutyl-imidazole (9.9 g) in DMF (25 ml). The reaction mixture was stirred overnight and quenched with H<sub>2</sub>O. The reaction mixture was filtered through celite and evaporated to yield an oily product. TLC showed a mixture of isomers.

**EXAMPLE 46****2'-O-(Imidazol-1-yl)butylguanosine**

A mixture of the 2,6-diamino-9-[2'-O-(imidazol-1-yl)butyl- $\beta$ -D-ribofuranosyl]purine and 2,6-diamino-9-[3'-O-(imidazol-1-yl)butyl- $\beta$ -D-ribofuranosyl]purine isomers in 0.1 M tris buffer (pH 7.4), 0.1 M NaSO<sub>4</sub> buffer (pH 7.4) and DMSO is treated with adenosine deaminase at RT for 5 days as per the procedure of Example 3. The product containing fractions are purified by silica gel chromatography and the product containing fraction evaporated to give the product.

**EXAMPLE 47****N2-Isobutyryl-2'-O-(imidazol-1-yl)butylguanosine**

2'-O-(imidazol-1-yl)butylguanosine in pyridine will be treated with trimethylsilyl chloride (5 eq) and isobutyryl chloride (5 eq) as per the procedure of Example 4 to yield the product.

**EXAMPLE 48****N2-Isobutyryl-5'-dimethoxytrityl-2'-O-(imidazol-1-yl)butylguanosine**

N2-Isobutyryl-2'-O-(imidazol-1-yl)butylguanosine will be treated with dimethoxytrityl chloride (1.1 eq), and dimethylaminopyridine (as a catalyst) in pyridine as per the procedure of Example 5. After chromatography purification, the product containing fractions will be evaporated to yield the product).

**EXAMPLE 49****2',3'-O-Dibutylstannylene uridine**

Utilizing the protocol of Wagner, et al., *J. Org. Chem.* 1974, 39, 24, uridine (45 g, 0.184 mol) was refluxed with  
5 di-n-butyltin oxide (45 g, 0.181 mol) in 1.4 l of anhydrous methanol for 4 hrs. The solvent was filtered and the resultant 2',3'-O-dibutylstannylene-uridine was dried under vacuum at 100°C for 4 hrs to yield 81 g (93%).

**EXAMPLE 50****10 2'-O-[Pentyl- $\omega$ -(N-phthalimido)]uridine**

2',3'-O-Dibutyl stannylene-uridine was dried over P<sub>2</sub>O<sub>5</sub> under vacuum for 12 hrs. To a solution of this compound (20 g, 42.1 mmols) in 500 ml of anhydrous DMF were added 25 g (84.2 mmols) of N(5-bromopentyl)phthalimide (Trans World Chemicals,  
15 Rockville, Maryland) and 12.75 g (85 mmols) of cesium fluoride (CsF). The mixture was stirred at room temperature for 72 hrs. The reaction mixture was evaporated then co-evaporated once with toluene and the residue was partitioned between EtOAc and water (400 ml each). The EtOAc layer was concentrated and  
20 applied to a silica column (700 g). Elution with CH<sub>2</sub>Cl<sub>2</sub>-CH<sub>3</sub>OH (20:1, v/v) gave fractions containing a mixture of the 2'- and 3'- isomers of O-pentyl- $\omega$ -N-phthalimido uridine, in 50% yield.

**EXAMPLE 51****5'-O-Dimethoxytrityl-2'-O-[pentyl- $\omega$ -(N-phthalimido)]uridine**

25 The isomeric mixture of 2'-O-[pentyl- $\omega$ -(N-phthalimido)]uridine was allowed to react with DMT chloride in dry pyridine at room temperature for 6 hrs. CH<sub>3</sub>OH was used to quench excess DMT-Cl and the residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub> containing 0.5% Et<sub>3</sub>N and water. The organic layer was  
30 dried (MgSO<sub>4</sub>) and the residue was applied to a silica column. The column was eluted with CH<sub>2</sub>Cl<sub>2</sub>:CH<sub>3</sub>OH (20:1, v/v) to separate the 2' and 3' isomers of the product.

- 49 -

**EXAMPLE 52**

**5'-O-Dimethoxytrityl-2'-O-[pentyl- $\omega$ -(N-phthalimido)]uridine-3'-O-( $\beta$ -cyanoethyl N,N-diisopropylphosphoramidite)**

5'-O-Dimethoxytrityl-2'-O-[pentyl- $\omega$ -(N-phthalimido)]uridine was placed in a dry round bottom flask containing a teflon stir-bar. The flask was purged with argon. Anhydrous methylene chloride was added to the flask in an amount sufficient to dissolve the nucleoside. Previously vacuum dried N,N-diisopropylaminohydrotetrazolide (600 mg, 0.014 mol) was added under argon. Bis-N,N-diisopropylamino-cyanoethylphosphite was added via syringe. The reaction was stirred under argon at 25°C for 16 h. Upon completion of the reaction, the reaction was transferred to a separatory funnel. The reaction flask was rinsed with methylene chloride (2x50 mL). The combined organic layer was washed 2 x with sat'd aq. sodium bicarbonate. The organic layer was dried over magnesium sulfate, evaporated and taken up in toluene containing 1% triethylamine. The resulting phosphoramidite was purified by silica gel flash chromatography and eluted with 3:1  $\rightarrow$  1:1 Hexanes/ethyl acetate containing 1% triethylamine. Selected fractions were combined, concentrated under reduced pressure and dried to yield the product as a white foam.  $^{31}\text{P}$ -NMR ( $\text{CDCl}_3$ ,  $\text{H}_3\text{PO}_4$  std.) showed the correct diastereomers

**EXAMPLE 53****25 2'-O-Pentyluridine**

Utilizing the procedures of Examples 50 and 51, 2',3'-O-dibutylstannylene uridine (19.1 g) was treated with bromopentane (7 ml, 1.3 eq.) and sodium iodide (4.5 g) in DMF (90 ml). Purification on a silica gel column utilizing MeOH/ $\text{CH}_2\text{Cl}_2$  5%  $\rightarrow$  10% yielded the a mixture of 2' and 3' isomers of the product as a dark oil (9.8 g).

**EXAMPLE 54****5'-O-Dimethoxytrityl-2'-O-pentyluridine**

The mixture of 2'-O-pentyluridine and 3'-O-pentyluridine (9.8 g) was reacted with dimethoxytrityl chloride

- 50 -

(10.5 g) as per the procedure of Example 51. The crude product was purified on a silica gel column (1000 g). Elution with Hex.-EtOAc (3:1 → 1:1) gave 5.5 g of the 2'-O-pentyl isomer and 3 g of the 3'-O-pentyl isomer. Anal. Calcd. for  $C_{35}H_{37}N_2O_8 \cdot \frac{1}{2}H_2O$ :  
5 C, 67.51; H, 6.55; N, 4.5. Found: C, 67.48; H, 6.55; N, 4.5.

**EXAMPLE 55****5'-O-Dimethoxytrityl-2'-O-pentyluridine-3'-O-(β-cyanoethyl N,N-diisopropylphosphoramidite)**

The protected 5'-O-dimethoxytrityl-2'-O-pentyluridine  
10 (4.6 g, 0.007 mol) was placed in a dry round bottom flask containing a teflon stir-bar. The flask was purged with argon. Anhydrous methylene chloride was added to the flask in an amount sufficient to dissolve the nucleoside. Previously  
vacuum dried N,N-diisopropylaminohydroxytetrazolide (600 mg,  
15 0.014 mol) was added under argon. Bis-N,N-diisopropylamino-cyanoethylphosphite (4.5 g, 4.7 ml, 2 eq.) was added with stirring via syringe. The reaction was stirred under argon at 25°C for 16 h. After verifying the completion of the reaction by TLC, the reaction was transferred to a separatory funnel and  
20 the reaction flask was rinsed with methylene chloride (2x50 mL). The combined organic layer was washed 2 x with sat'd aq. sodium bicarbonate. The organic layer was dried over magnesium sulfate, evaporated and taken up in toluene containing 1% triethylamine. The resulting phosphoramidite was purified by  
25 silica gel flash chromatography (300 g) and eluted with Hexanes/ethyl acetate (3:1 → 1:1 containing 1% triethylamine). Selected fractions were combined, concentrated under reduced pressure and dried to yield 2.67 g of product as a white foam.  
 $^{31}P$ -NMR ( $CDCl_3$ ,  $H_3PO_4$  std.) showed the correct diastereomers

**30 EXAMPLE 56****2'-O-Methyluridine**

As per the procedure of Example 49, uridine (8.5 g) was treated with dibutyl tin oxide (8.2 g, 1 eq). The resulting 2',3'-O-dibutylstannylene uridine was treated with

- 51 -

iodomethane (16 ml) at 42°C as per Example 50 to give a mixture of the 2' and 3' alkylated products (3.5 g) as a foam.

**EXAMPLE 57****5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-methyluridine**

5           2'-O-Methyluridine (8.0 g, 0.031 mol) was evaporated under reduced pressure with pyridine (100 mL) to an oil. To the residue was added 4,4'-dimethoxytriphenylmethyl chloride (DMT-Cl, 11.5 g, 0.34 mol) and pyridine (100 mL). The mixture was stirred at 25°C for 1.5 h and then quenched by the addition  
10 of methanol (10 mL) for 30 min. The mixture was concentrated under reduced pressure and the residue was chromatographed on silica gel (250 g), Elution with hexanes-ethyl acetate-triethylamine (80:20:1) and then ethyl acetate-triethylamine (99:1). The appropriate fractions were combined, evaporated  
15 under reduced pressure and dried at 25°C/0.2 mmHg for 1 h to provide 17.4 g (100 %) of tan foam; TLC purity 98% (R<sub>f</sub> 0.23, hexanes-ethyl acetate 4:1); PMR (DMSO) δ 11.4 (H-N<sup>3</sup>), 7.78 (H-6), 7.6-6.8 (Bz), 5.8 (H-1'), 5.3 (H-5'), 5.25 (HO-3'), 3.7 (CH<sub>3</sub>O-Bz), 3.4, (CH<sub>3</sub>O-2').

**20 EXAMPLE 58****5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-methyluridine-3'-O-(β-cyanoethyl N,N-diisopropylphosphoramidite)**

The product was prepared as per Example 54 from the intermediate 5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-methyl-  
25 uridine. Ethyl acetate-hexanes-triethylamine (59:40:1) was used as the chromatography eluent to give the product as a solid foam in 60% yield. TLC homogenous diastereomers, R<sub>f</sub> 0.58; 0.44 [ethyl acetate-hexanes-triethylamine 59:40:1)]. <sup>31</sup>P-NMR (CDCl<sub>3</sub>, H<sub>3</sub>PO<sub>4</sub> std.) δ 148.11; 148.61 (diastereomers).

**30 EXAMPLE 59****2'-O-Propyluridine**

As per the procedure of Example 49, uridine (10 g) was treated with dibutyl tin oxide (10.2 g, 1 eq). The resulting 2',3'-O-dibutylstannylene uridine was treated with iodopropane

- 52 -

(8 ml, 2 eq.) at 110°C as per Example 50 to give a mixture of the 2' and 3' isomers (5.5 g) as a foam.

**EXAMPLE 60****5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-propyluridine**

5           The mixture of 2'-O-propyluridine and 3'-O-propyluridine (3.6 g) was reacted with dimethoxytrityl chloride (4.2 g, 1.0 eq.) as per Example 51. The residue was chromatographed on silica gel eluted with Hex/EtOAc (1:1 with 1% triethylamine). The appropriate fractions were combined, 10 evaporated under reduced pressure and dried to provide 4.2 g of a white foam. Anal. Calcd. for  $C_{33}H_{36}N_2O_8 \cdot \frac{1}{2}H_2O$ : C, 67.33; H, 6.16; N, 4.76. Found: C, 67.15; H, 6.24; N, 4.44.

**EXAMPLE 61****5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-propyluridine-3'-O-  
15 (β-cyanoethyl N,N-diisopropylphosphoramidite)**

The product was prepared as per Example 54 from the intermediate 5'-O-(4,4'-dimethoxytrityl)-2'-O-propyluridine (470 mg) to yield the product as a foam (477 mg).

**EXAMPLE 62****20 2'-O-Nonyluridine**

As per the procedure of Example 49, uridine (22.5 g) was treated with dibutyl tin oxide (22.5 g, 1 eq). The resulting 2',3'-O-dibutylstannyleneuridine was treated with iodononane (11 ml, 1.3 eq.) at 130-140°C as per Example 50 to 25 give the 2' and 3' isomers (11.2 g) as an oil.

**EXAMPLE 63****5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-nonyluridine**

The mixture of 2'-O-nonyluridine and 3'-O-nonyluridine (11.2 g) was reacted with dimethoxytrityl chloride (10.5 g) as 30 per Example 51. The residue was chromatographed on silica gel eluted with Hex/EtOAc (3:1 → 1:1 with 1% triethylamine). The appropriate fractions were combined, evaporated under reduced pressure and dried to provide 5.2 g of a foam. An analytical

- 53 -

sample was rechromatographed using toluene/EtOAc (3:1 with 1% triethylamine) Anal. Calcd. for  $C_{39}H_{48}N_2O_8$ : C, 69.62; H, 7.19; N, 4.16. Found: C, 69.66; H, 7.18; N, 4.06.

**EXAMPLE 64**

- 5 **5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-nonyluridine-3'-O-( $\beta$ -cyanoethyl N,N-diisopropylphosphoramidite)**

The product was prepared as per Example 54 from the intermediate 5'-O-(4,4'-dimethoxytrityl)-2'-O-nonyluridine (3.1 g) to yield the product as a foam (2.49 g).

10 **EXAMPLE 65**

**2'-O-Hexenyluridine**

- As per the procedure of Example 49, uridine (10.5 g) was treated with dibutyl tin oxide (10.5 g, 1 eq). The resulting 2',3'-O-dibutylstannyleneuridine was treated with 6-  
15 bromohexene (3.5 ml, 1.2 eq.) and sodium iodide (3.3 g, 1.0 eq.) at 115°C as per Example 50 to give the 2' and 3' isomers (3.3 g) as a foam.

**EXAMPLE 66****5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-hexenyluridine**

- 20 The mixture of 2'-O-hexenyluridine and 3'-O-hexenyluridine (3.1 g) was reacted with dimethoxytrityl chloride (3.5 g, 1.1 eq.) as per Example 51. The residue was chromatographed on silica gel eluted with Hex/EtOAc (3:1  $\rightarrow$  1:1 with 1% triethylamine). The appropriate fractions were  
25 combined, evaporated under reduced pressure and dried to provide 2.3 g of a white foam. Anal. Calcd. for  $C_{36}H_{40}N_2O_8 \cdot \frac{1}{2}H_2O$ : C, 67.80; H, 6.48; N, 4.39. Found: C, 68.77; H, 6.41; N, 4.45.

**EXAMPLE 67**

- 30 **5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-hexenyluridine-3'-O-( $\beta$ -cyanoethyl N,N-diisopropylphosphoramidite)**

The product is prepared as per Example 54 from the intermediate 5'-O-(4,4'-dimethoxytrityl)-2'-O-hexenyluridine.

- 54 -

**EXAMPLE 68**

5'-O-Dimethoxytrityl-2'-O-[hexyl- $\omega$ -(N-phthalimido)]uridine-3'-O-( $\beta$ -cyanoethyl N,N-diisopropylphosphoramidite)

In a like manner as per Examples 50 through 52, using N-(6-bromohexyl)phthalimide, an N-phthalimide substituted hexyl group was introduced at the 2'-position of uridine followed by dimethoxytritylation and phosphitylation to give the title nucleotide.

**EXAMPLE 69**

10 5'-O-Dimethoxytrityl-2'-O-[decyl- $\omega$ -(N-phthalimido)]uridine-3'-O-( $\beta$ -cyanoethyl N,N-diisopropylphosphoramidite)

In a like manner as per Examples 50 through 52, using N-(10-bromodecyl)phthalimide, an N-phthalimide substituted decyl group was introduced at the 2'-position of uridine followed by dimethoxytritylation and phosphitylation to give the title nucleotide.

**EXAMPLE 70**

**N4-Benzoyl-2'-O-methylcytidine, Method A**

Step 1. 3',5'-O-[(1,1,3,3-Tetraisopropyl)-1,3-disiloxanediyl]cytidine

20 With stirring, cytidine (40 g, 0.165 mol) and 1,3-dichloro-1,1,3,3-tetraisopropylidisiloxane (TIPS-Cl, 50 g, 0.159 mol) were added to dry pyridine (250 mL). After stirring for 16 h at 25°C, the reaction was concentrated under reduced pressure to an oil. The oil was dissolved in methylene chloride (800 mL) and washed with sat'd sodium bicarbonate (2x300 mL). The organic layer was passed through a silica gel (200 g) scrub column. The product was recovered by elution with methylene chloride-methanol (97:3). The appropriate fractions were combined, evaporated under reduced pressure and dried at 25°C/0.2 mmHg for 1 h to give 59.3 g (77%) of oil. TLC purity 95% (Rf 0.59, ethyl acetate-methanol 9:1). The product may be crystallized from ethyl acetate as white crystals, mp 242-244°C. 41 PMR (DMSO)  $\delta$  7.7 (H-6), 5.68 (H-5), 5.61 (HO-2'), 5.55 (H-1').



- 55 -

**Step 2. N4-Benzoyl-3'-5'-O-[(1,1,3,3)tetraisopropyl-1,3-disiloxanediyl]cytidine**

Benzoyl chloride (18.5 g, 0.13 mol) was added over 30 min to a stirred solution of 3',5'-O-[(1,1,3,3-tetraisopropyl)-1,3-disiloxanediyl]cytidine (58 g, 0.12 mol) and triethylamine (15.6 g, 0.16 mol) in dimethylacetamide (400 mL) at 5°C. The mixture was allowed to warm to 25°C for 16 h and then poured onto ice water (3.5 L) with stirring. The resulting solid was collected, washed with ice water (3x500 mL) and dried at 45°C/0.2 mmHg for 5 h to provide 77 g (100%) of solid. TLC purity ca. 90% (Rf 0.63, chloroform-methanol 9:1); PMR (CDCl<sub>3</sub>) d 8.32 (H-6); mp 100-101°C.

**Step 3. N4-Benzoyl-2'-O-methyl-3',5'-O-[(1,1,3,3)tetraisopropyl-1,3-disiloxanediyl]cytidine**

A mixture of N4-benzoyl-3'-5'-O-[(1,1,3,3)tetraisopropyl-1,3-disiloxanediyl]cytidine (166 g, 0.25 mol, 90% purity), silver oxide (150 g, 0.65 mol) and toluene (300 mL) was evaporated under reduced pressure. More toluene (500 mL) was added and an additional amount (100 mL) was evaporated. Under a nitrogen atmosphere, methyl iodide was added in one portion and the reaction was stirred at 40°C for 16 h. The silver salts were collected and washed with ethyl acetate (3x150 mL). The combined filtrate was concentrated under reduced pressure. The residue was dissolved in a minimum of methylene chloride, applied to a silica gel column (1 kg) and eluted with hexanes-ethyl acetate (3:2 → 1:1). The appropriate fractions were combined, concentrated under reduced pressure and dried at 45°C/0.2 mmHg for 1 h to yield 111 g (66%) of oil; TLC purity ca. 90% (Rf 0.59, hexanes-ethyl acetate 3:2). PMR (CDCl<sub>3</sub>) d 8.8 (br s, 1, H-N<sup>4</sup>), 8.40 (d, 1, H-6), 8.0-7.4 (m, 6, H-5 and Bz), 5.86 (s, 1, H-1'), 3.74 (s, 3, CH<sub>3</sub>O-2').

**Step 4. N4-Benzoyl-2'-O-methylcytidine**

A solution of N4-benzoyl-2'-O-methyl-3',5'-O-[(1,1,3,3)tetraisopropyl-1,3-disiloxanediyl]cytidine (111 g, 0.18 mol) in methanol (160 mL) and tetrahydrofuran (640 mL) was treated with tetrabutylammonium fluoride solution (368 mL, 1 M

- 56 -

in tetrahydrofuran). The reaction was stirred at 25°C for 16 h. The pH was adjusted to 7 with Amberlite IRC-50 resin. The mixture was filtered and the resin was washed with hot methanol (2x200 mL). The combined filtrate was concentrated under  
5 reduced pressure, absorbed on silica gel (175 g) and chromatographed on silica gel (500 g, ethyl acetate-methanol 19:1 → 4:1). Selected fractions were combined, concentrated under reduced pressure and dried at 40°C/0.2 mmHg for 2 h to yield 28 g (42.4%, 21.5% from cytidine) of solid; TLC  
10 homogenous (R<sub>f</sub> 0.37, ethyl acetate). mp 178-180°C (recryst. from ethanol); PMR (CDCl<sub>3</sub>) δ 11.22 (br s, 1, H-N<sup>4</sup>), 8.55 (d, 1, H-6), 8.1-7.2 (m, 6, H-5 and Bz), 5.89 (d, 1, H-1'), 5.2 (m, 2, HO-3',5'), 3.48 (s, 3, CH<sub>3</sub>O-2').

**EXAMPLE 71****15 N<sup>4</sup>-Benzoyl-2'-O-methylcytidine, Method B****Step 1. 2'-O-methylcytidine**

Cytidine (100 g, 0.41 mol) was dissolved in warm dimethylformamide (65°C, 1125 mL). The solution was cooled with stirring to 0°C. A slow, steady stream of nitrogen gas was  
20 delivered throughout the reaction. Sodium hydride (60% in oil, washed thrice with hexanes, 18 g, 0.45 mol) was added and the mixture was stirred at 0°C for 45 min. A solution of methyl iodide (92.25 g, 40.5 mL, 0.65 mol) in dimethylformamide (400 mL) was added in portions over 4 h at 0°C. The mixture was  
25 stirred for 7 h at 25°C and then filtered. The filtrate was concentrated to dryness under reduced pressure followed by co-evaporation with methanol (2x200 mL). The residue was dissolved in methanol (350 mL). The solution was adsorbed on silica gel (175 g) and evaporated to dryness. The mixture was  
30 slurried in dichloromethane (500 mL) and applied on top of a silica gel column (1 kg). The column was eluted with a gradient of dichloromethane-methanol (10:1 → 2:1). The less polar 2',3'-dimethyl side product was removed and the co-eluting 2' and 3'-O-methyl product containing fractions were  
35 combined and evaporated under reduced pressure to a syrup. The syrup was dissolved in a minimum of hot ethanol (ca. 150 mL)

- 57 -

and allowed to cool to 25°C. The resulting precipitate (2' less soluble) was collected, washed with ethanol (2x25 ml) and dried to give 15.2 g of pure 2'-O-methylcytidine; mp 252-254°C mp 252-254°C); TLC homogenous (Rf 0.50, dichloromethane-methanol 3:1, (Rf of 3' isomer 0.50 and the dimethyl product 0.80). The filtrate was evaporated to give 18 g of a mixture of isomers and sodium iodide.

#### Step 2. N4-Benzoyl-2'-O-methylcytidine

The pure 2'-O-methylcytidine (15.2 g, 0.060 mol) was dissolved in a solution of benzoic anhydride (14.7 g, 0.12 mol) in dimethylformamide (200 mL). The solution was stirred at 25°C for 48 h and then evaporated to dryness under reduced pressure. The residue was triturated with methanol (2x200 mL), collected and then triturated with warm ether (300 mL) for 10 min. The solid was collected and triturated with hot 2-propanol (50 mL) and allowed to stand at 4°C for 16 h. The solid was collected and dried to give 17 g of product. The crude filtrate residue (18 g) of 2'-O-methylcytidine was treated with benzoic anhydride (17.3 g, 0.076 mol) in dimethylformamide (250 mL) as above and triturated in a similar fashion to give an additional 6.7 g of pure product for a total yield of 23.7 g (16% from cytidine) of solid; TLC homogenous (Rf 0.25, chloroform-methanol 5:1, co-spots with material made utilizing Method A)

#### 25 EXAMPLE 72

##### N4-Benzoyl-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-methylcytidine

N4-Benzoyl-2'-O-methylcytidine, (28 g, 0.077 mol) was evaporated under reduced pressure to an oil with pyridine (400 mL). To the residue was added 4,4'-dimethoxytriphenylmethyl chloride (DMT-Cl, 28.8 g, 0.085 mol) and pyridine (400 mL). The mixture was stirred at 25°C for 2 h and then quenched by the addition of methanol (10 mL) for 30 min. The mixture was concentrated under reduced pressure and the residue was chromatographed on silica gel (500 g, hexanes-ethyl acetate-triethylamine 60:40:1 and then ethyl acetate-triethylamine

- 58 -

99:1). The appropriate fractions were combined, evaporated under reduced pressure and dried at 40°C/0.2 mmHg for 2 h to give 26 g (74%) of foam; TLC homogenous (Rf 0.45, ethyl acetate); PMR (DMSO)  $\delta$  11.3 (H-N<sup>4</sup>), 8.4-6.9 (H-6, H-5, Bz), 5.95 (H-1'), 5.2 (HO-3'), 3.7 (s, 6, CH<sub>3</sub>O-trit.), 3.5 (s, 3, CH<sub>3</sub>O-2')

**EXAMPLE 73**

**N4-Benzoyl-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-methylcytidine-3'-O-( $\beta$ -cyanoethyl N,N-diisopropylphosphoramidite)**

10 The product was prepared as per the procedure of Example 38 starting with intermediate compound N4-benzoyl-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-methylcytidine (22.0 g, 0.0333 mole) and using ethyl acetate-hexanes-triethylamine (59:40:1) as the chromatography eluent to give the product as  
15 a solid foam (23.6 g) in 83% yield; TLC homogenous diastereomers (Rf 0.46; 0.33, ethyl acetate-hexanes-triethylamine 59:40:1); <sup>31</sup>P-NMR (CD<sub>3</sub>CN, H<sub>3</sub>PO<sub>4</sub> std.)  $\delta$  150.34; 151.02 (diastereomers).

**EXAMPLE 74****20 2'-O-Nonylcytidine**

Cytidine (10.1 g, 0.0415 mol), sodium hydride (2.0 g, 1.2 eq), iodononane (9.8 ml, 1.2 eq.) in DMF (100 ml) were reacted as per the procedure of Example 71, Step 1 to yield the 2' and 3' isomers as a sticky foam (11.6 g).

**25 EXAMPLE 75****N4-Benzoyl-2'-O-nonylcytidine**

The mixture of 2'-O-nonylcytidine and 3'-O-nonylcytidine (11.5 g) is converted to N4-benzoyl-2'-O-nonylcytidine as per the procedure of Example 71, Step 2.

- 59 -

**EXAMPLE 76****N4-Benzoyl-5'-O-(dimethoxytrityl)-2'-O-nonylcytidine**

N4-Benzoyl-2'-O-nonylcytidine (2.67 g, 0.0056 mol) was treated with dimethoxytrityl chloride (2.0 g, 1.1 eq) as per the procedure of Example 72 to give 4.2 g of pure product. Anal. Calcd. for  $C_{46}H_{53}N_3O_8 \cdot \frac{1}{2}H_2O$ : C, 70.39; H, 6.93; N, 5.35. Found: C, 71.20; H, 6.88; N, 5.41.

**EXAMPLE 77****N4-Benzoyl-5'-O-(dimethoxytrityl)-2'-O-nonylcytidine-3'-O-( $\beta$ -cyanoethyl N,N-diisopropylphosphoramidite)**

The product was prepared as per the procedure of Example 38 starting with intermediate compound N4-benzoyl-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-nonylcytidine (4.1 g, 0.0053 mole) treated with bis-N,N-diisopropylaminocycanoethylphosphite (3.3 ml) and N,N-diisopropylaminohydroxytetrazolide (450 mg). The product was eluted from the silica gel column using Hexane/EtOAc (3:1  $\rightarrow$  1:1 with 1% triethylamine) as the chromatography eluent to give the product as a solid foam (4.21 g).  $^{31}P$ -NMR ( $CD_3CN$ ,  $H_3PO_4$  std.) shows the diastereomers;

**EXAMPLE 78****2'-O-Pentylcytidine**

Cytidine (10 g, 0.041 mol), sodium hydride (2.4 g, 1.5 eq), bromopentane (7.6 ml, 1.5 eq.) in DMSO (90 ml) were reacted as per the procedure of Example 71, Step 1 to yield the 2' and 3' isomers as a foam (7.6 g).

**EXAMPLE 79****N4-Benzoyl-2'-O-pentylcytidine**

The mixture of 2'-O-pentylcytidine and 3'-O-pentylcytidine (7.5 g) is converted to N4-benzoyl-2'-O-pentylcytidine as per the procedure of Example 71, Step 2.

- 60 -

**EXAMPLE 80****N4-Benzoyl-5'-O-(dimethoxytrityl)-2'-O-pentylcytidine**

N4-Benzoyl-2'-O-pentylcytidine (3.0 g, 0.007 mol) was treated with dimethoxytrityl chloride (2.7 g, 1.1 eq) as per the procedure of Example 72 to give 3.5 g of pure product. Anal. Calcd. for  $C_{42}H_{45}N_3O_8 \cdot \frac{1}{2}H_2O$ : C, 69.21; H, 6.36; N, 5.76. Found: C, 69.51; H, 6.30; N, 5.71.

**EXAMPLE 81****N4-Benzoyl-5'-O-(dimethoxytrityl)-2'-O-pentylcytidine-3'-O-( $\beta$ -cyanoethyl N,N-diisopropylphosphoramidite)**

The product was prepared as per the procedure of Example 38 starting with intermediate compound N4-benzoyl-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-pentylcytidine (3.5 g, 0.0048 mole) treated with bis-N,N-diisopropylaminocycanoethylphosphite (2.9 g, 3.1 ml, 2 eq.) and N,N-diisopropylaminohydroxytetrazolide (400 mg, 0.5 eq.). The product was eluted from the silica gel column using Hexane/EtOAc (3:1  $\rightarrow$  1:1 with 1% triethylamine) as the chromatography eluent to give the product as a solid foam (3.24 g).  $^{31}P$ -NMR ( $CD_3CN$ ,  $H_3PO_4$  std.) shows the diastereomers.

**EXAMPLE 82****2'-O-Propylcytidine**

Cytidine (16.5 g, 0.068 mol) was treated with sodium hydride (4.5 g) and bromopropane (15 ml) in DMF (150 ml) at room temperature for three days. The resulting reaction mixture was used directly in the next step (see Example 83).

**EXAMPLE 83****N4-Benzoyl-2'-O-propylcytidine**

To the 2'-O-propylcytidine reaction mixture of Example 82 in an ice bath was added pyridine (60 ml) and trimethylsilyl chloride (60 ml). The reaction was stirred for 30 mins followed by the addition of benzoyl chloride (55 ml). The resulting reaction mixture was stirred for 2.5 hrs and then cooled in an ice bath.  $H_2O$  (100 ml) and conc.  $NH_4OH$  (100 ml)

- 61 -

were added. After stirring for 30 mins, the reaction mixture was evaporated and the residue partition between  $H_2O$  and  $CH_2Cl_2$ . The organic phase was washed once with dil  $Na_2CO_3$ , once with dil  $HCl$ , dried over  $MgSO_4$  and evaporated. The resulting residue  
5 was loaded on a silica gel column (150 g) and eluted with first  $CH_2Cl_2$  then 5 to 10%  $MeOH$  in  $CH_2Cl_2$  as the elution solvent. The product containing fractions were evaporated to a foam. The foam was crystallized from  $EtOAc/Hexanes$  to give the product (6.5 g total) in several crystal batches.

**10 EXAMPLE 84****N4-Benzoyl-5'-O-(dimethoxytrityl)-2'-O-propylcytidine**

N4-Benzoyl-2'-O-propylcytidine (3.0 g, 0.007 mol) was treated with dimethoxytrityl chloride (1.5 g) as per the procedure of Example 72 to give 1.5 g of pure product. Anal.  
15 Calcd. for  $C_{40}H_{42}N_3O_8 \cdot \frac{1}{2}H_2O$ : C, 68.45; H, 6.18; N, 5.99. Found: C, 68.39; H, 5.99; N, 5.95.

**EXAMPLE 85****N4-Benzoyl-5'-O-(dimethoxytrityl)-2'-O-propylcytidine-3'-O-( $\beta$ -cyanoethyl N,N-diisopropylphosphoramidite)**

20 The product was prepared as per the procedure of Example 38 starting with intermediate compound N4-benzoyl-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-O-propylcytidine (3.8 g, 0.0055 mole) treated with bis-N,N-diisopropylaminocycanoethylphosphite (3.5 ml, 2 eq.) and N,N-  
25 diisopropylaminohydroxotetrazolide (500 mg, 0.5 eq.). The product was eluted from the silica gel column using Hexane/ $EtOAc$  (1:1 with 1% triethylamine) as the chromatography eluent to give the product as a solid foam (4.72 g).  $^{31}P$ -NMR ( $CD_3CN$ ,  $H_3PO_4$  std.) shows the diastereomers.

**30 EXAMPLE 86****N2,N6-Diisobutyrylamino-9-(2'-O-methyl- $\beta$ -D-ribofuranosyl)purine**

N2,N6-Diamino-9-(2'-O-methyl- $\beta$ -D-ribofuranosyl)purine (1.6 g, 5.39 mmol, see Example 34) was co-evaporated with pyridine (25 ml). A suspension of the residue in pyridine (40

- 62 -

ml) was cooled in an ice bath and trimethylsilyl chloride (4.8 ml) was added. The reaction mixture was stirred for 30 mins followed by the addition of butyryl chloride (2.8 ml, 5 eq). The resulting reaction mixture was stirred at room temperature  
5 for 4 hours. H<sub>2</sub>O (10 ml) and conc. NH<sub>4</sub>OH (10 ml) were added with stirring to quench the reaction mixture. After 30 mins, the reaction mixture was evaporated and the residue purified on a silica gel column using CH<sub>2</sub>Cl<sub>2</sub> → 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to elute the product. The appropriate fractions were evaporated to yield  
10 the product as an oil (2.4 g).

**EXAMPLE 87****N2,N6-Diisobutyrylamino-9-(5'-O-dimethoxytrityl-2'-O-methyl-β-D-ribofuranosyl)purine**

N2,N6-Diisobutyrylamino-9-(2'-O-methyl-β-D-ribofuranosyl)purine (2.4 g) was co-evaporated with pyridine and redissolved in pyridine. Dimethoxytrityl chloride (1.8 g, 1 eq) and dimethylaminopyridine (5 mg) were added and the resulting solution was stirred overnight at room temperature. The solvent was partly evaporated and the residue partition  
15 between CH<sub>2</sub>Cl<sub>2</sub> - dil. Na<sub>2</sub>CO<sub>3</sub>. The organic phase was washed with dil. Na<sub>2</sub>CO<sub>3</sub>, dried with MgSO<sub>4</sub> and evaporated. The residue was purified on a silica gel column eluted with Hexanes/EtOAc (1:1) containing 1% triethylamine. The fraction contain the product were evaporated to yield the product as a foam (2.4 g).

**25 EXAMPLE 88****N2,N6-Diisobutyrylamino-9-[5'-O-dimethoxytrityl-2'-O-methyl-3'-O-(β-cyanoethyl N,N-diisopropylphosphoramidate)-β-D-ribofuranosyl]purine**

N2,N6-Diisobutyrylamino-9-(5'-O-dimethoxytrityl-2'-O-methyl-β-D-ribofuranosyl)purine (1.7 g, 0.0023 mol) was treated  
30 with bis-N,N-diisopropylaminocynoethylphosphite (1.48 ml, 2 eq.) and N,N-diisopropylaminohydrotetrazolide (200 mg) at room temperature overnight. The reaction mixture was partitioned between dil. Na<sub>2</sub>CO<sub>3</sub>/CH<sub>2</sub>Cl<sub>2</sub>, the organic phase was dried over  
35 MgSO<sub>4</sub> and evaporated. The residue was loaded on a silica gel



- 63 -

column and eluted with Hexanes/EtOAc (3:1 → 1:1 with 1% triethylamine) to give the product as a solid foam (1.73 g). <sup>31</sup>P-NMR (CD<sub>3</sub>CN, H<sub>3</sub>PO<sub>4</sub> std.) shows the diastereomers.

**EXAMPLE 89****5 Oligonucleotide Synthesis**

Once nucleoside phosphoramidites of the invention have been prepared, they can then subsequently be incorporated into oligonucleotides, which are synthesized by a standard solid phase, automated nucleic acid synthesizer such as the Applied Biosystems, Incorporated 380B or MilliGen/Bioscience 7500 or 8800. Standard phosphoramidite coupling chemistries (see, e.g., M. Caruthers, *Oligonucleotides. Antisense Inhibitors of Gene Expression.*, pp. 7-24, J.S. Cohen, ed., CRC Press, Inc. Boca Raton, FL, 1989) are used with these synthesizers to provide the desired oligonucleotides. The Beaucage reagent (see, e.g., *J. Am. Chem. Soc.* 1990, 112, 1253) or elemental sulfur (see, e.g., *Tetrahedron Letters* 1981, 22, 1859), can likewise be used to provide phosphorothioate oligonucleotides.

**EXAMPLE 90****20 A. Evaluation of the thermodynamics of hybridization of 2'- modified oligonucleotides.**

The ability of the 2'- modified oligonucleotides to hybridize to their complementary RNA or DNA sequences is determined by thermal melting analysis. The RNA complement is synthesized from T7 RNA polymerase and a template-promoter of DNA synthesized with an Applied Biosystems, Inc. 380B RNA species was purified by ion exchange using FPLC (LKB Pharmacia, Inc.). Natural antisense oligonucleotides or those containing 2'-O-alkyl guanosine at specific locations are added to either the RNA or DNA complement at stoichiometric concentrations and the absorbance (260 nm) hyperchromicity upon duplex to random coil transition was monitored using a Gilford Response II spectrophotometer. These measurements are performed in a buffer of 10 mM Na-phosphate, pH 7.4, 0.1 mM EDTA, and NaCl to yield an ionic strength of 10 either 0.1 M or 1.0 M.

- 64 -

Data is analyzed by a graphic representation of  $1/T_m$  vs  $\ln[Ct]$ , where  $[Ct]$  was the total oligonucleotide concentration. From this analysis the thermodynamic parameters is determined. Based upon the information gained concerning the stability of the duplex of heteroduplex formed, the placement of 2'-O-alkyl guanosine into oligonucleotides are assessed for their effects on helix stability. Modifications that drastically alter the stability of the hybrid exhibit reductions in the free energy ( $\Delta G$ ) and decisions concerning their usefulness as antisense oligonucleotides are made.

#### **B. Fidelity of hybridization of 2'- modified oligonucleotides**

The ability of the 2'-O-alkyl guanosine modified antisense oligonucleotides to hybridize with absolute specificity to the targeted mRNA is shown by Northern blot analysis of purified target mRNA in the presence of total cellular RNA. Target mRNA is synthesized from a vector containing the cDNA for the target mRNA located downstream from a T7 RNA polymerase promoter. Synthesized mRNA was electrophoresed in an agarose gel and transferred to a suitable support membrane (ie. nitrocellulose). The support membrane was blocked and probed using [ $^{32}P$ ]-labeled antisense oligonucleotides. The stringency will be determined by replicate blots and washing in either elevated temperatures or decreased ionic strength of the wash buffer. Autoradiography was performed to assess the presence of heteroduplex formation and the autoradiogram quantitated by laser densitometry (LKB Pharmacia, Inc.). The specificity of hybrid formation was determined by isolation of total cellular RNA by standard techniques and its analysis by agarose electrophoresis, membrane transfer and probing with the labeled 2'-modified oligonucleotides. Stringency was predetermined for the unmodified antisense oligonucleotides and the conditions used such that only the specifically targeted mRNA was capable of forming a heteroduplex with the 2'-modified oligonucleotide.

- 65 -

**EXAMPLE 91 - Nuclease Resistance****A. Evaluation of the resistance of 2'- modified oligonucleotides to serum and cytoplasmic nucleases.**

Natural phosphorothioate, and 2- modified oligo-  
5 nucleotides were assessed for their resistance to serum nucleases by incubation of the oligonucleotides in media containing various concentrations of fetal calf serum or adult human serum. Labeled oligonucleotides were incubated for various times, treated with protease K and then analyzed by gel  
10 electrophoresis on 20% polyacrylamine-urea denaturing gels and subsequent autoradiography. Autoradiograms were quantitated by laser densitometry. Based upon the location of the modifications and the known length of the oligonucleotide it was possible to determine the effect on nuclease degradation by  
15 the particular 2'-modification. For the cytoplasmic nucleases, a HL60 cell line was used. A post-mitochondrial supernatant was prepared by differential centrifugation and the labeled oligonucleotides were incubated in this supernatant for various times. Following the incubation, oligo-nucleotides were  
20 assessed for degradation as outlined above for serum nucleolytic degradation. Autoradiography results were quantitated for comparison of the unmodified, the phosphorothioates, and the 2'- modified oligonucleotides.

**25 B. Evaluation of the resistance of 2'- modified oligonucleotides to specific endo- and exo-nucleases.**

Evaluation of the resistance of natural and 2'-modified oligonucleotides to specific nucleases (ie, endonucleases, 3',5'-exo-, and 5',3'-exonucleases) was done to determine the exact effect of the modifications on degradation.  
30 Modified oligonucleotides were incubated in defined reaction buffers specific for various selected nucleases. Following treatment of the products with proteinase K, urea was added and analysis on 20% poly- acrylamide gels containing urea was done. Gel products were visualized by staining using Stains All  
35 (Sigma Chemical Co.). Laser densitometry was used to quantitate


- 66 -

the extend of degradation. The effects of the 2'-modifications were determined for specific nucleases and compared with the results obtained from the serum and cytoplasmic systems.

- 67 -

What is claimed is:

1. A compound having the structure:



wherein X is  $R_1-(R_2)_n$ ;

$R_1$  is  $C_3$ - $C_{20}$  alkyl,  $C_4$ - $C_{20}$  alkenyl or  $C_2$ - $C_{20}$  alkynyl;


$R_2$  is halogen, hydroxyl, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, N-phthalimido, imidazole, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, polyamine, polyamide, polyalkylene glycol, polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, or a group that enhances the pharmacokinetic properties of oligonucleotides; and

n is an integer from 0 to about 6.

2. The compound of claim 1 wherein  $R_1$  is  $C_4$ - $C_{20}$  alkyl.
3. The compound of claim 1 wherein  $R_1$  is  $C_5$ - $C_{20}$  alkyl.

- 68 -

4. A compound having the structure:



wherein X is  $R_1-(R_2)_n$ ;

$R_1$  is  $C_3-C_{20}$  alkyl;

$R_2$  is  $NH_2$ , imidazole, or N-phthalimido;

Y is a hydroxyl blocking group;

Z is phosphate or an activated phosphate group;

$Q_1$  and  $Q_2$  independently are H or a guanosine blocking group; and

n is an integer from 0 to about 6.


5. The compound of claim 4 wherein:

Y is trityl, methoxytrityl, dimethoxytrityl or trimethoxytrityl.

6. The compound of claim 4 wherein:

Z is  $\beta$ -cyanoethyl-N,N-isopropylphosphoramidate.

7. A compound having the structure:



wherein X is  $R_1-(R_2)_n$ ;

$R_1$  is  $C_3-C_{20}$  alkyl,  $C_4-C_{20}$  alkenyl or  $C_2-C_{20}$  alkynyl;

$R_2$  is halogen, hydroxyl, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl,

- 69 -

NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, imidazole, N-phthalimido, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, polyamine, polyamide, polyalkylene glycol, polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, or a group that enhances the pharmacokinetic properties of oligonucleotides; and

n is an integer from 0 to about 6.

8. The compound 2'-O-propylguanosine, 2'-O-pentylguanosine, 2'-O-nonylguanosine, 2'-O-octadecylguanosine, 2'-O-(N-phthalimido)-pentylguanosine, or 2'-O-(imidazol-1-yl)butylguanosine.

9. A process for preparing a 2'-O-alkylated guanosine 3'-O-phosphoramidite comprising the steps of:

alkylating a 2,6-diamino-9-(ribofuranosyl)purine to form a 2,6-diamino-9-(2'-O-alkylated ribofuranosyl)purine;

deaminating said 2,6-diamino-9-(2'-O-alkylated ribofuranosyl)purine to form a 2'-O-alkylated guanosine;

blocking the 5'-hydroxyl moiety of said 2'-O-alkylated guanosine; and

phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated guanosine.

10. The process of claim 9 wherein said alkylation is in the presence of base.

11. The process of claim 10 wherein said base is a metal hydride.

12. The process of claim 9 wherein said blocking step adds a dimethoxytrityl group.

- 70 -

13. The process of claim 9 wherein said phosphitylation step is conducted with bis-N,N-diisopropyl-aminocyanoethylphosphite.

14. The process of claim 13 wherein said phosphitylation step is conducted in the presence of N,N-diisopropylamino-hydrotetrazolide.

15. The process of claim 9 further comprising blocking the N2-amino moiety of said 2'-O-alkylated guanosine prior to blocking said 5' hydroxyl moiety.

16. The process of claim 9 wherein said deamination step is performed using adenosine deaminase.

17. A process for preparing 2'-O-alkyl-guanosine comprising:

treating 2,6-diaminopurine riboside with a base and a compound having the formula R-L wherein R is said alkyl group and L is a leaving group, to form 2'-O-alkyl-2,6-diaminopurine riboside and 3'-O-alkyl-2,6-diaminopurine riboside; and

deaminating said 2'-O-alkyl-2,6-diaminopurine riboside to yield the said 2'-O-alkyl guanosine.

18. The process of claim 17 further comprising:  
separating said 2'-O-alkyl-2,6-diaminopurine riboside from said 3'-O-alkyl-2,6-diaminopurine riboside.

19. The process of claim 18 wherein said 2'-O-alkyl-2,6-diaminopurine riboside is separated from said 3'-O-alkyl-2,6-diaminopurine riboside by conducting said deamination under conditions such that the rate of deamination of said 2'-O-alkyl-2,6-diaminopurine riboside is accelerated compared to the rate of deamination of said 3'-O-alkyl-2,6-diaminopurine riboside.



- 71 -

20. The process of claim 17 wherein said 2'-O-alkyl-2,6-diaminopurine riboside is deaminated with adenosine deaminase.

21. The process of claim 19 wherein said accelerated rate of deamination of said 2'-O-alkyl-2,6-diaminopurine riboside is achieved by use of the enzyme adenosine deaminase.

22. The process of claim 17 wherein said base is sodium hydride.

23. The process of claim 17 wherein said 2'-O-alkyl-2,6-diaminopurine riboside is physically separated from said 3'-O-alkyl-2,6-diaminopurine riboside by crystallization or chromatography.

24. The process of claim 17 wherein said alkyl group is C<sub>1</sub>-C<sub>20</sub> straight chain or branched chain, saturated or unsaturated alkyl, C<sub>1</sub>-C<sub>20</sub> alicyclic, C<sub>1</sub>-C<sub>20</sub> straight chain or branched chain, saturated or unsaturated, substituted alkyl or C<sub>1</sub>-C<sub>20</sub> substituted alicyclic.

25. The process of claim 24 wherein the substituent of said C<sub>1</sub>-C<sub>20</sub> straight chain or branched chain, saturated or unsaturated, substituted alkyl or C<sub>1</sub>-C<sub>20</sub> substituted alicyclic is halogen, hydroxyl, thiol, keto, carboxyl, nitrates), nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, heterocyclic, alicyclic, carbocyclic, intercalators, reporter molecules, conjugates, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligonucleotides, and groups that enhance the pharmacokinetic properties of oligonucleotides.

- 72 -

26. A process for preparing 2'-O-alkyl-guanosine comprising:

treating 2,6-diaminopurine riboside with a base and a compound having the formula R-L wherein R is said alkyl group and L is a leaving group, to form 2'-O-alkyl-2,6-diaminopurine riboside; and

treating said 2'-O-alkyl-2,6-diaminopurine riboside with adenosine deaminase.

27. The process of claim 26 wherein said alkyl group is a C<sub>1</sub>-C<sub>20</sub> straight chain or branched chain, saturated or unsaturated alkyl or a C<sub>1</sub>-C<sub>20</sub> straight chain or branched chain, saturated or unsaturated, substituted alkyl.

28. The process of claim 27 wherein the substituent of said C<sub>1</sub>-C<sub>20</sub> straight chain or branched chain, saturated or unsaturated, substituted alkyl is halogen, hydroxyl, thiol, keto, carboxyl, nitrates), nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, heterocyclic, alicyclic, carbocyclic, intercalators, reporter molecules, conjugates, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligonucleotides, and groups that enhance the pharmacokinetic properties of oligonucleotides.

29. The process of claim 26 wherein said base is sodium hydride.

30. The process of claim 26 further comprising:  
separating said 2'-O-alkyl-2,6-diaminopurine riboside from further alkylated products resulting from said treatment of said 2,6,-diaminopurine riboside with said base and said compound having the formula R-L wherein R is said alkyl group and L is a leaving group.

- 73 -

31. The process of claim 27 wherein said alkyl group is C<sub>1</sub>-C<sub>20</sub> saturated, straight chain alkyl or C<sub>1</sub>-C<sub>20</sub> saturated, straight chain substituted alkyl.

32. A process for preparing a 3'-O-phosphoramidite of 2'-O-alkyl guanosine comprising:

treating 2,6-diaminopurine riboside with a base and a compound having the formula R-L wherein R is said alkyl group and L is a leaving group, to form 2'-O-alkyl-2,6-diaminopurine riboside;

treating said 2'-O-alkyl-2,6-diaminopurine riboside with adenosine deaminase to form 2'-O-alkyl-guanosine;

protecting the 2-NH<sub>2</sub> moiety of said 2'-O-alkyl-guanosine with a protecting group;

protecting the 5'-OH moiety of said 2'-O-alkyl-guanosine with a further protecting group; and

phosphitylating the 3'-OH moiety of said protected 2'-O-alkyl-guanosine.

33. The process of claim 32 wherein said protected 2'-O-alkyl guanosine is phosphitylated with 2-cyanoethyl N,N-diisopropylaminochlorophosphine.

34. The process of claim 32 wherein said base is sodium hydride.

35. A process for preparing 2'-O-alkyl-2,6-diaminopurine riboside, 3'-O-alkyl-2,6-diaminopurine riboside and 2',3'-di-O-alkyl-2,6-diaminopurine riboside comprising:

treating 2,6-diaminopurine riboside with a base and a compound having the formula R-L wherein R is said alkyl group and L is a leaving group, to form at least one of 2'-O-alkyl-2,6-diaminopurine riboside, 3'-O-alkyl-2,6-diaminopurine riboside or 2',3'-di-O-alkyl-2,6-diaminopurine riboside; and

- 74 -

isolating one of said 2'-O-alkyl-2,6-diaminopurine riboside, 3'-O-alkyl-2,6-diaminopurine riboside or 2',3'-di-O-alkyl-2,6-diaminopurine riboside.

36. The process of claim 35 further comprising:

adding sufficient compound having the formula R-L wherein R is said alkyl group and L is a leaving group, and base to said 2,6-diaminopurine riboside to form a mixture of at least two of said 2'-O-alkyl-2,6-diaminopurine riboside, 3'-O-alkyl-2,6-diaminopurine riboside and 2',3'-di-O-alkyl-2,6-diaminopurine riboside; and

separating out said least one of said 2'-O-alkyl-2,6-diaminopurine riboside, 3'-O-alkyl-2,6-diaminopurine riboside and 2',3'-di-O-alkyl-2,6-diaminopurine riboside from said mixture.

37. The process of claim 36 further comprising:

separating out said at least one of said 2'-O-alkyl-2,6-diaminopurine riboside, 3'-O-alkyl-2,6-diaminopurine riboside or 2',3'-di-O-alkyl-2,6-diaminopurine riboside from said mixture via crystallization.

38. The process of claim 36 further comprising:

separating out said at least one of said 2'-O-alkyl-2,6-diaminopurine riboside, 3'-O-alkyl-2,6-diaminopurine riboside or 2',3'-di-O-alkyl-2,6-diaminopurine riboside from said mixture via chromatography.

39. A process for preparing the 3'-O-phosphoramidite of 2,6-diamino-9-(2'-O-alkyl- $\beta$ -D-ribofuranosyl)purine comprising:

treating 2,6-diaminopurine riboside with a base and a compound having the formula R-L wherein R is said alkyl group and L is a leaving group to form 2'-O-alkyl-2,6-diaminopurine riboside;

- 75 -

protecting the 2 and 6 -NH<sub>2</sub> moieties of said 2,6-diamino-9-(2'-O-alkyl-β-D-ribofuranosyl)purine with protecting groups;

protecting the 5'-OH moiety of said 2,6-diamino-9-(2'-O-alkyl-β-D-ribofuranosyl)purine with a further protecting group; and

phosphitylating the 3'-OH moiety of said protected 2,6-diamino-9-(2'-O-alkyl-β-D-ribofuranosyl)purine.

40. An improved process for the preparation of 2,6-diaminopurine riboside comprising:

treating guanosine hydrate with hexamethyldisilazane and trifluoromethane sulfonic acid in a sealed container with the exclusion of liquid ammonia and under non-anhydrous conditions; and

isolating said 2,6-diaminopurine riboside by crystallization.

41. A process for preparing a 2'-O-alkylated cytidine 3'-O-phosphoramidite comprising the steps of:

alkylating an unblocked cytidine to form a 2'-O-alkylated cytidine;

blocking the 5'-hydroxyl moiety of said 2'-O-alkylated cytidine; and

phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated cytidine.

42. The process of claim 41 wherein said alkylation is in the presence of base.

43. The process of claim 42 wherein said base is a metal hydride.

44. The process of claim 41 wherein said blocking step adds a dimethoxytrityl group.

- 76 -

45. The process of claim 41 wherein said phosphitylation step is conducted with bis-N,N-diisopropylaminocynoethylphosphite.

46. The process of claim 45 wherein said phosphitylation step is conducted in the presence of N,N-diisopropylamino-hydrotetrazolide.

47. The process of claim 41 further comprising blocking the N4-amino moiety of said 2'-O-alkylated cytidine prior to blocking said 5' hydroxyl moiety.

48. A process for preparing a 2'-O-alkylated uridine 3'-O-phosphoramidite comprising the steps of:

treating a uridine with a dialkyltin oxide to form a 2',3'-O-dialkylstannylene derivative of uridine;

alkylating said stannylene derivative of uridine to form a 2'-O-alkylated uridine;

blocking the 5'-hydroxyl moiety of said 2'-O-alkylated uridine; and

phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated uridine.

49. The process of claim 48 wherein said alkylation is in the presence of a salt.

50. The process of claim 48 wherein said salt is a metal halide.

51. The process of claim 48 wherein said blocking step adds a dimethoxytrityl group.

52. The process of claim 48 wherein said phosphitylation step is conducted with bis-N,N-diisopropylaminocynoethylphosphite.

- 77 -

53. The process of claim 52 wherein said phosphitylation step is conducted in the presence of N,N-diisopropylamino-hydrotetrazolide.

54. The process of claim 48 wherein said dialkyl tin oxide is dibutyl tin oxide.

55. A process for preparing a 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine 3'-O-phosphoramidite comprising the steps of:

alkylating a 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine to provide a 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine;

blocking the 5'-hydroxyl moiety of said 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine; and

phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine.

56. The process of claim 55 wherein said alkylation is in the presence of a base.

57. The process of claim 56 wherein said base is a metal hydride.

58. The process of claim 55 wherein said blocking step adds a dimethoxytrityl group.

59. The process of claim 55 wherein said phosphitylation step is conducted with bis-N,N-diisopropylaminocynoethylphosphite.


60. The process of claim 59 wherein said phosphitylation step is conducted in the presence of N,N-diisopropylamino-hydrotetrazolide.

61. The process of claim 55 further comprising blocking the N2 and N6-amino moieties of said 2'-O-alkylated

- 78 -

2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine prior to blocking said 5' hydroxyl moiety.

62. An oligomer comprising at least one subunit having the structure:



wherein X is  $R_1-(R_2)_n$ ;

$R_1$  is  $C_3$ - $C_{20}$  alkyl,  $C_4$ - $C_{20}$  alkenyl, or  $C_2$ - $C_{20}$  alkynyl;  
 $R_2$  is halogen, hydroxyl, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, imidazole, N-phthalimido, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, polyamine, polyamide, polyalkylene glycol, polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, or a group that enhances the pharmacokinetic properties of oligonucleotides;


$T_3$  and  $T_5$  independently are OH or a further subunit of said oligomer that is joined to said structure; and

n is an integer from 0 to about 6.



- 79 -

63. An oligomer comprising at least one subunit having the structure:



wherein X is  $R_1-(R_2)_n$ ;

$R_1$  is  $C_1$ - $C_{20}$  alkyl,  $C_2$ - $C_{20}$  alkenyl, or  $C_2$ - $C_{20}$  alkynyl;  
 $R_2$  is halogen, hydroxyl, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, imidazole, N-phthalimido, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, polyamine, polyamide, polyalkylene glycol, polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, or a group that enhances the pharmacokinetic properties of oligonucleotides;

$T_3$  and  $T_5$  independently are OH or a further subunit of said oligomer that is joined to said structure; and

$n$  is an integer from 0 to about 6.

64. A process for preparing an oligonucleotide that comprises at least one 2'-O-alkylated guanosine nucleotide comprising:

alkylating a 2,6-diamino-9-(ribofuranosyl)purine to form a 2,6-diamino-9-(2'-O-alkylated ribofuranosyl)purine;

deaminating said 2,6-diamino-9-(2'-O-alkylated ribofuranosyl)purine to form a 2'-O-alkylated guanosine;

- 80 -

blocking the 5'-hydroxyl moiety of said 2'-O-alkylated guanosine;

phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated guanosine to form a 2'-O-alkylated guanosine 3'-O-phosphoramidite; and

coupling said 2'-O-alkylated guanosine 3'-O-phosphoramidite with a 5'-hydroxyl moiety on an oligonucleotide utilizing phosphoramidite coupling conditions.

65. A process for preparing an oligonucleotide that comprises at least one 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine nucleotide within the sequence of the oligonucleotide comprising:


alkylating a 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine to provide a 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)-purine;

blocking the 5'-hydroxyl moiety of said 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine;

phosphitylating the 3'-position of said 5'-blocked 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine to form a 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine 3'-O-phosphoramidite; and

coupling said 2'-O-alkylated 2,6-diamino-9-( $\beta$ -D-ribofuranosyl)purine 3'-O-phosphoramidite to a 5'-hydroxyl moiety on an oligonucleotide utilizing phosphoramidite chemistry.

66. A method of modulating the synthesis of a protein comprising specifically hybridizing with mRNA coding for said protein an oligomer comprising at least one subunit having the structure:



- 81 -

wherein X is  $R_1-(R_2)_n$ ;

$R_1$  is  $C_3-C_{20}$  alkyl,  $C_4-C_{20}$  alkenyl, or  $C_2-C_{20}$  alkynyl;


$R_2$  is halogen, hydroxyl, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, imidazole, N-phthalimido, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, polyamine, polyamide, polyalkylene glycol, polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, and a group that enhances the pharmacokinetic properties of oligonucleotides;

$T_3$  and  $T_5$  independently are OH or a further nucleotide or nucleoside of said oligonucleotide or oligonucleoside that is joined to said structure; and

n is an integer from 0 to about 6.

67. The method of claim 66 wherein said oligonucleotide is in a pharmaceutically acceptable carrier.

68. A method of modulating the synthesis of a protein comprising specifically hybridizing with mRNA coding for said protein an oligomer comprising at least one subunit having the structure:



wherein X is  $R_1-(R_2)_n$ ;

- 82 -

$R_1$  is  $C_1$ - $C_{20}$  alkyl,  $C_2$ - $C_{20}$  alkenyl, or  $C_2$ - $C_{20}$  alkynyl;  $R_2$  is halogen, hydroxyl, thiol, keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, amino, imidazole, N-phthalimido, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, aryl, heterocycle, carbocycle, intercalator, reporter molecule, conjugate, polyamine, polyamide, polyalkylene glycol, polyether, a group that enhances the pharmacodynamic properties of oligonucleotides, and a group that enhances the pharmacokinetic properties of oligonucleotides;

$T_3$  and  $T_5$  independently are OH or a further nucleotide or nucleoside of said oligonucleotide or oligonucleoside that is joined to said structure; and

$n$  is an integer from 0 to about 6.

69. The method of claim 68 wherein said oligonucleotide is in a pharmaceutically acceptable carrier.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/06807

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C07H 21/02; 19/067, 19/10, 19/173, 19/20; A61K 31/70

US CL : 536/24.5, 25.3, 26.8, 27.7, 27.81, 28.53, 28.54, 514/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/24.5, 25.3, 26.8, 27.7, 27.81, 28.53, 28.54, 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS AND CAS ONLINE,

SEARCH TERMS: 2'-O-ALKYL, 2'-O-METHYL, 2'-O-ETHYL, SODIUM HYDRIDE, NUCLEOSIDES, ALKYLATION

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Proceedings of the National Academy of Sciences, volume 87, issued October 1990, Iribarren et al., "2'-O-Alkyl oligonucleotides as antisense probes", pages 7747 - 7751, especially page 7747, column 2, first sentence above "Materials and Methods".	1 - 69
Y	Nucleic Acids Research, Volume 19, No. 4, issued 1991, Sproat et al., "New synthetic routes to synthons suitable for 2'-O-allyl-oligoribonucleotide assembly", pages 733 - 738, esp Abstract and page 733, column 1.	1 - 69

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be part of particular relevance		
"E" earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed	"&"	document member of the same patent family

Date of the actual completion of the international search

08 OCTOBER 1993

Date of mailing of the international search report

21 OCT 1993

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

GARY L. KUNZ

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US93/06807

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Nucleic Acids Research, Vol. 19, No. 10, Cotten et al., "2'-O-methyl, 2'-O-ethyl oligonucleotides and phosphorothioate oligodeoxyribonucleotides as inhibitors of the in vitro UR snRNP-dependent mRNA processing event", pages 2629 - 2635, esp. Abstract on page 2629.	1 - 69
Y	Nucleic Acids Research, Vol. 19, No. 21, issued 1991, Wagner et al., "A simple procedure for the preparation of protected 2'-O-methyl or 2'-O-ethyl ribonucleoside-3'-phosphoramidites", pages 5965 - 5971, esp. page 5965, Abstract.	1 - 69
Y	Nucleic Acids Research, Vol. 18, No. 1, issued 1990, Sproat et al., "New synthetic routes to protected purine 2'-O-methylriboside-3'-O-phosphoramidites using a novel alkylation procedure", pages 41 - 49, esp. page 41, columns 1 - 2.	1 - 69
Y	Nucleic Acids Research, Vol. 15, Number 15, 1987, Inoue et al., "Synthesis and hybridization studies on two complementary nona(2'-O-methyl)ribonucleotides", pages 6131 - 6148, esp. page 6131, Abstract.	1 - 69
Y	Canadian Journal of Chemistry, Volume 59; issued 1981, Robins et al., "Nucleic Acid related compounds. 36. Synthesis of the 2'-O-methyl and 3'-O-methyl ethers of guanosine and 2'-aminoadenosine and correlation of O'-methylnucleoside <sup>13</sup> C nmr specral shifts", pages 3360 - 3364, see esp. the Reaction Scheme on page 3361.	1 - 69